

RESEARCH ARTICLE

Open Access

Theoretical study on the electronic, structural, properties and reactivity of a series of mono-, di-, tri- and tetrachlorothiophenes as well as corresponding radical cation forms as monomers for conducting polymers

Hossein Shirani IL Beigi^{1*}, Saeed Jameh-Bozorghi²

Abstract

In this paper, electrical and structural properties of mono-, di-, tri- and tetrachlorothiophenes and their radical cations have been studied using the density functional theory and B3LYP method with 6-311++G** basis set. The effects of the number and position of the substituent of chlorine atoms on the properties of the thiophene ring for all chlorothiophenes and their radical cations have been studied. Vibrational frequencies, nuclear chemical shielding constants, spin-density distribution, size and direction of dipole moment vector, ionization potential, electric polarizabilities and NICS values of these compounds have been calculated as well. The analysis of these data showed that double bonds in 3-chlorothiophene are more delocalized and it is the best possible candidate monomer among all chlorothiophenes for the synthesis of corresponding conducting polymers with modified characteristics.

Introduction

Electrically-conducting polymers such as advanced materials and the new class of 'synthetic metals' have drawn high attention in the last years [1]. The 2000 Nobel Prize in chemistry recognized the discovery of conducting polymers and over 25 years of progress in this field [2,3]. There has been growing interest in research on conducting polymer nano structures (nanorods, -tubes, -wires and -fibers) since they incorporate the advantages of organic conductors with low-dimensional systems and so create interesting physicochemical properties and potentially useful applications [4-7]. The varieties of such studies have been reported in the literature that includes: surfactants [8-11], liquid crystals [12], polyelectrolytes [13], nano-wire seeds [14], thiophene oligomers [15], bulky organic dopants [16-21]. Among intrinsic conducting polymers with conjugated double

bonds, polythiophenes (PTHs) and related derivatives are suitable for commercial applications [22].

Polythiophenes are normally produced from non aqueous media because the monomer is more soluble in them. The influence of water on polymerization process of thiophene as well as on the redox switching properties has been studied [23,24]. It has been recently shown that the attachment of a range of electron-donating or electron-withdrawing groups through a conjugated linker has a dramatic effect on polymerization potential and subsequent photovoltaic performance [25,26]. Polythiophene has a solubility of 75% (w/w) in chloroform and a conductivity of 220 s cm^{-1} [22].

The effect of temperature on polymerization process of thiophene has been investigated as well [27]. Results show that when polymerization is carried out at 15-20°C, polymers with optimal properties will be obtained. Ultrasonication has been used to improve the efficiency (improved yield, lowering of polymerization potentials) of electropolymerization process of polythiophene [28].

* Correspondence: shiranihossein@gmail.com

¹Young Researchers Club, Islamic Azad University, Toyserkan Branch, Toyserkan, Iran

Full list of author information is available at the end of the article


Thiophene is oxidized to form a conducting electro active polymer (CEP), with the best conductivity when it is created from α - α linkages [22]. In polypyrroles, the mechanism of polymerization involves the formation of radical cations which react with another monomer to develop the polymeric structure (Figure 1) [29].

The objective of the present research is to study electronic and structural properties of all chlorothiophenes (shown in Figure 2) and their radical cations using DFT-B3LYP and DFT-UB3LYP methods.

The electropolymerization and characteristics of monomers (conductivity and solubility) are related to the stability of their radical cations which it is closely dependent on the type and configuration of the substituent on the thiophene ring [22]. Vibrational frequencies with analyses of their IR intensities and NMR shielding constants will be used to characterize and predict the molecular and spectroscopic properties of fluorothiophenes.

Computational Procedures

First of all, structures of representative mono-, di-, tri- and tetrachlorothiophenes were optimized using B3LYP/6-311++G** method [30-32]. The geometries of these compounds were optimized using standard gradient techniques with default parameters set in G98W. Vibrational frequencies with their IR intensities are calculated with scale factors, $f_1=1.0100$ for low-frequency vibrations (below 1000 cm^{-1}) and $f_2=0.9679$ for high-frequency vibrations (above 1000 cm^{-1}) [33]. The vibrational



frequencies analysis showed that all structures correspond to local minima in potential energy surface.

Results and Discussion

Structural Analysis of Chlorothiophenes


In this section, the electrical and structural properties of chlorothiophenes and their related radical cations have been studied. Because of the stabilizing effects of the delocalization of non-bonding p electrons of sulfur atom, it is expected that the thiophene molecules possess a plane structure.

In agreement with this expectation, results of the present calculations show that all chlorothiophenes possess planar structures (the optimized values of all dihedral angles of all structures are either 0° or 180°). The optimized structural parameters obtained at B3LYP/6-311++G** level of theory for all chlorothiophenes and UB3LYP/6-311++G** level of theory for all radical cations are listed in Tables 1 and 2.

To predict the bonding-characteristic behavior of chlorothiophene rings in their corresponding polymer chains and to determine the extent of the π -conjugation character of these polymers, we have used the F_n coefficient defined in Equation 1 for each thiophene ring [34].

$$F_n = \frac{R34}{(R23 + R45)/2} \quad (1)$$

where R34 is the length of C_3 - C_4 bond (formally single bond) and $[(R23 + R45)/2]$ is the average length of C_2 - C_3 and C_4 - C_5 bonds (formally double bonds) in the thiophene ring. The calculated values of the F_n coefficient for thiophene, all chlorothiophenes and their radical cations are listed in Table 1. As shown in this table, F_n coefficient has the following trend for all molecules:



And for all molecular radical cations, values of the F_n coefficient has the following trend:



As this table shows, 3-chlorothiophene has the smallest value of the F_n coefficient. Therefore, it can be




Table 1 B3LYP/6-311++G optimized values of bond lengths (Å) and the F_n coefficient for Thiophene and Chlorothiophenes**

Molecule	S1-C2	S1-C5	C2-C3	C3-C4	C4-C5	F_n
Thiophene	1.733	1.733	1.366	1.428	1.366	1.045
	<i>1.728</i>	<i>1.728</i>	<i>1.422</i>	<i>1.376</i>	<i>1.421</i>	<i>0.968</i>
A	1.740	1.737	1.364	1.428	1.364	1.047
	<i>1.760</i>	<i>1.715</i>	<i>1.419</i>	<i>1.378</i>	<i>1.414</i>	<i>0.973</i>
B	1.732	1.733	1.363	1.424	1.364	1.044
	<i>1.681</i>	<i>1.780</i>	<i>1.437</i>	<i>1.341</i>	<i>1.366</i>	0.957
C	1.741	1.735	1.367	1.427	1.362	1.046
	<i>1.732</i>	<i>1.743</i>	<i>1.445</i>	<i>1.395</i>	<i>1.383</i>	<i>0.987</i>
D	1.740	1.737	1.362	1.427	1.361	1.048
	<i>1.782</i>	<i>1.700</i>	<i>1.385</i>	<i>1.386</i>	<i>1.431</i>	<i>0.984</i>
E	1.745	1.745	1.362	1.429	1.361	1.050
	<i>1.750</i>	<i>1.750</i>	<i>1.410</i>	<i>1.379</i>	<i>1.410</i>	<i>0.978</i>
F	1.730	1.730	1.363	1.433	1.363	1.051
	<i>1.733</i>	<i>1.734</i>	<i>1.413</i>	<i>1.387</i>	<i>1.412</i>	<i>0.982</i>
G	1.738	1.734	1.367	1.435	1.360	1.052
	<i>1.754</i>	<i>1.728</i>	<i>1.415</i>	<i>1.386</i>	<i>1.411</i>	<i>0.981</i>
H	1.747	1.744	1.365	1.429	1.360	1.049
	<i>1.734</i>	<i>1.768</i>	<i>1.430</i>	<i>1.383</i>	<i>1.390</i>	<i>0.981</i>
I	1.745	1.743	1.364	1.437	1.364	1.053
	<i>1.752</i>	<i>1.752</i>	<i>1.413</i>	<i>1.388</i>	<i>1.413</i>	<i>0.982</i>

Corresponding values for the related radical cations optimized with UB3LYP/6-311++G** are given in the lower row with italic fonts. See Figure 2 for definitions of bond lengths.

suggested that double bonds in 3-chlorothiophene are more delocalized.

Spin-Distribution Analysis

Electropolymerization process for each molecular radical cation is obtained from α - α (C_2 - C_5) linkages [29], so it can be said that in a compound, the greater spin density in Positions C_3 and C_4 is, the higher electropolymerization rate will be [35].

The spin-density distribution over the ring atoms of chlorothiophene radical cations have been calculated and presented in Table 3. As shown in this table, it is obvious that monomers **B** and **F** have the maximal spin density on positions C_3 and C_4 . This means that the electropolymerization of 3-chlorothiophene (**B**) and 3,4-dichlorothiophene (**F**) can be more favorable ratio with respect to other molecules.

Electric Dipole Moments

Orientations of the polymer chains in the condensed phase are the most important parameters affecting the electric charge transport properties of polymers so that the electrical conductivity of a polymer chain is altered when its orientation and consequently its symmetry and non isotropic interactions are changed.

Chemically synthesized PTHs powders exhibit a partially crystalline structure in which the chains align and

Table 2 B3LYP/6-311++G optimized values of bond angles for Thiophene and Chlorothiophenes**

Molecule	C2-S1-C5	S1-C2-W	S1-C5-X	S1-C2-C3	S1-C5-C4	C3-C4-C5	C2-C3-Y	C5-C4-Z
Thiophene	91.5	119.9	119.9	111.5	111.5	112.7	123.4	123.4
	<i>89.7</i>	<i>120.1</i>	<i>120.1</i>	<i>113.3</i>	<i>113.3</i>	<i>111.9</i>	<i>122.6</i>	<i>122.6</i>
A	90.7	120.5	119.6	112.5	111.8	111.8	123.4	123.3
	<i>88.0</i>	<i>120.0</i>	<i>119.4</i>	<i>114.9</i>	<i>114.2</i>	<i>112.8</i>	<i>122.7</i>	<i>122.3</i>
B	91.7	121.1	119.9	110.6	111.9	111.6	123.4	124.6
	<i>90.6</i>	<i>123.1</i>	<i>118.8</i>	<i>111.4</i>	<i>113.4</i>	<i>110.0</i>	<i>120.7</i>	<i>125.2</i>
C	91.0	120.9	120.8	119.6	112.2	112.3	124.2	124.5
	<i>88.2</i>	<i>122.9</i>	<i>118.2</i>	<i>113.4</i>	<i>115.4</i>	<i>110.7</i>	<i>120.6</i>	<i>124.5</i>
D	90.9	120.4	120.4	112.8	110.8	114.6	124.5	123.4
	<i>88.5</i>	<i>118.4</i>	<i>121.9</i>	<i>115.5</i>	<i>112.4</i>	<i>115.1</i>	<i>125.0</i>	<i>120.4</i>
E	89.9	120.1	120.0	112.7	112.7	112.4	123.4	123.4
	<i>86.3</i>	<i>119.4</i>	<i>119.4</i>	<i>115.8</i>	<i>115.8</i>	<i>111.1</i>	<i>122.5</i>	<i>122.5</i>
F	91.9	120.9	120.9	111.3	111.3	112.8	123.6	123.6
	<i>89.9</i>	<i>122.9</i>	<i>121.5</i>	<i>113.0</i>	<i>112.9</i>	<i>112.1</i>	<i>123.1</i>	<i>123.2</i>
G	91.3	120.6	120.5	112.1	111.5	113.4	124.4	123.5
	<i>88.3</i>	<i>121.2</i>	<i>120.7</i>	<i>114.5</i>	<i>114.1</i>	<i>112.9</i>	<i>123.2</i>	<i>122.8</i>
H	90.3	120.1	120.4	113.0	111.6	113.5	124.5	124.2
	<i>86.7</i>	<i>121.8</i>	<i>118.0</i>	<i>114.1</i>	<i>116.8</i>	<i>109.4</i>	<i>120.9</i>	<i>124.5</i>
I	90.6	120.1	120.1	112.2	112.2	112.5	124.2	124.2
	<i>86.9</i>	<i>120.4</i>	<i>120.4</i>	<i>115.4</i>	<i>115.4</i>	<i>111.2</i>	<i>123.2</i>	<i>123.2</i>

Corresponding values for the related radical cations are given in the lower row with italic fonts. See Figure 2 for definitions of angles.

Table 3 Distribution of spin density over sulfur and carbon atoms in UB3LYP/6-311++G optimized structures of Thiophene and Chlorothiophenes radical cations**

Molecule	S ₁	C ₂	C ₃	C ₄	C ₅
Thiophene	-0.131	0.547	0.046	0.046	0.548
A	-0.087	0.377	0.055	0.049	0.456
B	0.041	0.356	0.289	-0.179	0.357
C	-0.081	0.358	0.186	-0.108	0.399
D	0.003	0.315	-0.124	0.162	0.425
E	-0.088	0.343	0.061	0.061	0.343
F	0.419	-0.110	0.236	0.240	-0.109
G	-0.065	0.384	-0.044	0.014	0.470
H	-0.062	0.349	0.133	-0.064	0.313
I	-0.067	0.362	-0.006	-0.005	0.361


pack in a herringbone arrangement (Figure 3) in the crystalline regions [36].

These overall interactions can be well expressed in terms of interactions between local dipole moments of monomers. The local alignment of monomers' dipole moments in solution with respect to the orientation of the polymer chain determines the electrochemical properties of the polymer formed on electrode surface.

In addition, the orientation of monomer in the double layer of the solution in an electropolymerization cell depends on both size and direction of the dipole moment vectors of both monomers and polymer chain [35].

It is necessary to study the dipole moments of these chlorothiophenes to be able to predict their electropolymerization properties. The size of the dipole moment vector and its components calculated for chlorothiophenes in this study are presented in Table 4.

An analysis of calculated dipole moments shows that the orientation of the dipole moment vector is toward sulfur atom for all chlorothiophenes. The size and direction of the dipole moment vector is more dependent on the position (symmetry) of substituent than its number.



It can also be seen in Table 4 that the dipole moments for all compounds have the following trend:

$$C > F > G > A > B > E > I > H > D > \text{Thiophene}$$

The size of the dipole moment vector for monomers **C** and **F** is greater than other compounds. Therefore, compounds **C** and **F** have a high solubility in polar solvents. These characteristics increase the efficiency of electrochemical polymerization processes of these monomers. Moreover it can be seen in Table 4 that thiophene has a low solubility in polar solvents.

Ionization Potentials

The first step in the electropolymerization of conducting polymers is formation of intermediate radical cations from monomers which can be considered as an ionization reaction [37].



where M and M^+ denote monomer and its radical cation respectively. So, stability of intermediate radical cations has an important role in the electropolymerization process.

Therefore, it is useful to calculate electronic energy difference between the neutral monomer (as the initial species) and positively charged monomer (as the intermediate). This energy difference is proportional to Ionization Potential (IP) of the monomer. Calculated values of IP for chlorothiophenes based on Equation 2 are presented in Table 4. The trend of electrochemical stability of these monomers is:

$$F > B > \text{Thiophene} > G > D > C > A > I > H > E$$

Since ionization potential energies are directly proportional to the electrochemical oxidation potentials of the compounds, it can be said that the electrochemical stability of monomers **F** and **B** are greater than other compounds.

Electric Polarizabilities

Values of Polarizabilities of compounds containing halogen are used to interpret light scattering and intensities of vibrational Raman spectra [34]. Values of exact electrical polarizabilities for all chlorothiophenes have been calculated and listed in Table 4. Based on this table, it can be claimed that with an increase in the number of chlorine atoms on the thiophene ring, polarization values will increase as well.

The zero values obtained for polarizabilities α_{xz} and α_{yz} are consequences of the planar structure. The small values of α_{xy} polarizabilities indicate that the delocalization strength of the π -system of the ring dominates the anisotropic effects of substituent for asymmetric chlorothiophenes.

Table 4 Electric dipole moments, polarizability tensor elements, and ionization potentials calculated at B3LYP/6-311++G level of theory for the optimized structures of Thiophene and Chlorothiophenes**

Molecule	Electric dipole moment (D)			Polarizabilities (Å)				Ionization potentials (ev)
	μ_x	μ_y	μ_{tot}	α_{xx}	α_{yy}	α_{zz}	α_{xy}	IP
Thiophene	0.0000	-0.5169	0.5169	67.093	73.851	41.017	0.012	8.75
A	1.2229	-1.1156	1.6553	83.379	90.474	45.543	-0.424	8.53
B	0.4885	-1.3293	1.4162	73.843	99.340	45.713	0.914	8.77
C	2.0522	-0.6022	2.1388	105.378	99.760	50.761	1.770	8.56
D	-0.0085	-0.6768	0.6768	118.479	89.726	50.349	5.704	8.58
E	-0.0014	-1.1668	1.1668	124.387	85.513	50.238	0.006	8.38
F	1.9947	-0.3062	2.0181	113.075	91.086	50.781	-3.497	9.00
G	0.2028	-1.8053	1.8166	123.767	114.854	56.216	6.267	8.66
H	-0.3414	-0.6487	0.7331	117.232	125.636	55.339	14.351	8.44
I	0.7756	-0.6432	1.0077	135.731	138.834	61.933	7.424	8.50

μ_z , α_{xz} , and α_{yz} are essentially zero for all compounds because of the planar structure.

Vibrational Spectra

The vibrational frequencies for all chlorothiophenes were calculated using optimized structures at B3LYP/6-311++G** level of theory. Calculated IR absorption frequencies and intensities for all chlorothiophenes are presented in Table 5. To facilitate the comparison the frequencies are divided into low-frequency (below 1000 cm^{-1}) and high-frequency groups (above 1000 cm^{-1}). According to analysis of frequencies listed in Table 5, it can be suggested that thermal decomposition has the following trend:

Thiophene > B > A > F > D > C > E > H > G > I

Thiophene and 3-chlorothiophene have the highest low-frequency values among all these compounds; this means that thiophene and 3-chlorothiophene have the largest force constants for their bending modes of vibration. Higher values of low frequencies obtained for thiophene and 3-chlorothiophene can be regarded as higher protection from thermal decomposition for these molecules.

The calculated Zero-Point vibrational Energy (ZPE) for thiophene and all chlorothiophenes are listed in Table 6. Having in mind the contents of the table it can be said that ZPE for thiophene is greater than other compounds, as a result the relative thermal stability of thiophene is higher than other compounds.

Table 5 B3LYP/6-311++G calculated IR absorption frequencies in cm^{-1} (intensities in km/mol) for Thiophene and Chlorothiophenes**

Molecule	Low - frequency range	High - frequency range
Thiophene	460(0.67) 577 (0.0) 619(0.37) 697(0.0) 727(143) 751(0.41) 842(26.9) 886(1.63) 894(0.30) 928(0.0)	1017(3.15) 1069(4.45) 1070(0.0) 1234(11.37) 1348(0.71) 1397(11.2) 1501(0.24) 3093(3.85) 3107(3.7) 3139(0.04) 3142(1.20)
A	208(0.72) 264(0.13) 438(4.5) 471(4.17) 566(0.36) 655(5.28) 655(83.78) 854(20.46) 920(0.13) 969(50.73)	1033(11.23) 1075(1.96) 1206(9.34) 1329(2.19) 1406(48.15) 1512(8.91) 3100(3.84) 3120(0.21) 3143(0.78)
B	210(0.51) 289(0.11) 426(4.34) 462(0.0) 611(15.8) 642(4.36) 682(3.5) 783(84.55) 812(1.54) 862(24.13) 886(0.01) 912(44.03)	1061(7.04) 1184(15.57) 1184(8.76) 1342(24.13) 1395(1.59) 1499(29.97) 3119(0.42) 3145(0.39) 3155(3.64)
C	158(0.12) 175(0.04) 240(1.51) 317(1.12) 418(1.45) 476(1.16) 497(5.84) 608(14.78) 680(6.04) 717(47.74) 815(0.53) 892(64.40) 897(1.62) 991(40.60)	1078(0.73) 1144(15.97) 1327(16.33) 1405(27.92) 1508(27.40) 3118(0.85) 3144(1.22)
D	159(0.05) 189(0.27) 247(0.01) 327(0.52) 393(1.20) 466(6.33) 471(1.12) 505(11.54) 687(9.9) 732(32.32) 833(35.74) 834(23.82) 883(42.25) 978(38.80)	1066(12.21) 1158(10.25) 1313(31.22) 1409(33.17) 1506(70.38) 3132(1.66) 3154(6.58)
E	117(0.01) 194(0.10) 273(0.0) 318(0.73) 362(1.33) 486(14.08) 527(19.39) 552(0.0) 672(1.37) 728(3.32) 799(46.01) 884(0.0) 982(7.20) 985(140.11)	1048(0.42) 1189(5.80) 1196(1) 1419(33.77) 1521(38.74) 3111(0.94) 3122(0.14)
F	170(0.0) 179(0.48) 242(1.90) 335(0.14) 454(4.75) 458(0.65) 467(1.14) 633(0.0) 692(20.02) 707(0.0) 800(67.63) 860(1.4) 863(24.2) 967(62.3)	1104(19.3) 1142(0.1) 1318(58.4) 1491(7.9) 1491(13.2) 3154(6.8) 3155(3.10)
G	94(0.04) 179(0.02) 187(0.11) 237(1.3) 241(0.19) 363(1.3) 367(0.57) 456(3.83) 481(0.14) 541(3) 658(2.6) 732(33.4) 775(46.6) 856(3.9) 928(71.1)	1009(3.9) 1120(16.22) 1297(79.65) 1410(17.9) 1498(37.1) 3155(7.6)
H	115(0.0) 178(0.0) 173(0.05) 187(0.0) 279(0.15) 355(1.5) 364(1.5) 420(1.8) 493(8.6) 582(21) 505(1.8) 695(3.1) 831(20.1) 853(61.7) 972(4.15) 999(85.6)	1123(14.1) 1289(20.6) 1421(13.9) 1516(86) 3130(4.67)
I	88(0.0) 114(0.02) 177(0.0) 181(0.05) 191(0.09) 236(0.91) 271(0.0) 489(4.6) 363(0.14) 364(1.4) 403(2.6) 456(3.17) 514(2.15) 657(0.0) 671(22.11) 809(45.32) 896(61.9) 972(168)	1025(28) 1270(87) 1424(1.24) 1505(75)

Table 6 Zero-Point vibrational energy in kcal/mol, calculated at B3LYP/6-311++G level of theory for the optimized structures of Thiophene and Chlorothiophenes**

Molecule	ZPE
Thiophene	41.64
A	35.85
B	35.81
C	30.01
D	29.92
E	29.97
F	29.93
G	24.07
H	24.09
I	18.20

NMR Chemical Shifts

The NMR chemical shifts can be used to predict ring current and estimate aromaticity of all chlorothiophenes [38-40]. ^{13}C chemical shielding have been calculated for DFT-B3LYP/6-311++G** optimized geometries of all chlorothiophenes. The isotropic and anisotropic magnetic shielding have been calculated using GIAO and CSGT methods [41-43].

Values of chemical shifts in Positions C_2 and C_3 for all chlorothiophenes are calculated and presented in Figure 4. It can be seen from this figure that compounds **B** and

F have the lowest chemical shifts in Position C_2 , these molecules have the highest chemical shifts in Position C_3 , as well. It should be noted that these compounds had the highest values of spin-density in their Position C_3 .

NICS Analysis

The Nuclear Independent Chemical Shift (NICS) at GIAO/B3LYP/6-311++G** and magnetic susceptibilities at CSGT/B3LYP/6-311++G** level of theory [44-49] in center, 0.5 and 1 angstrom upper than thiophene ring have been calculated and listed in Table 7. Negative values of NICS represent the magnetic shielding and electric current in cyclic compounds. NICS results showed that compounds **B** and **F** have the highest NICS values in rings center, 0.5 and 1 angstrom upper than thiophene ring; these compounds have the largest electrical current and most conductivity among all other compounds. Based on this table, it can be claimed that the magnetic susceptibilities of these molecules increase with the number of the chlorine atoms, as well.

Conclusion

B3LYP method with 6-311++G** basis set calculations have been carried out successfully to study the structural, electric and spectroscopic properties of all chlorothiophenes. Values of dipole moment, spin density distribution, ionization potential, vibrational frequencies




Figure 4 GIAO calculated values of the chemical shifts for Thiophene and chlorothiophenes (referenced to TMS); (a) $^{13}\text{C}_2$ chemical shifts, (b) $^{13}\text{C}_3$ chemical shifts.

Table 7 Calculated NICS values (ppm) at GIAO/B3LYP/6-311++G and magnetic susceptibilities (cgs-ppm) at CSGT/B3LYP/6-311++G** for Thiophene and Chlorothiophenes**

Molecule	NICS(0)	NICS(0.5)	NICS(1)	χ
Thiophene	-11.91	-11.49	-9.28	-51.86
A	-12.30	-11.54	-9.05	-68.66
B	-13.49	-12.22	-9.57	-69.26
C	-12.57	-11.45	-8.67	-86.79
D	-12.77	-11.50	-8.56	-85.45
E	-12.13	-10.91	-8.19	-85.24
F	-13.51	-12.12	-9.31	-86.81
G	-13.48	-11.87	-8.60	-104.38
H	-12.48	-10.90	-7.81	-103.01
I	-13.31	-11.38	-7.93	-123.23

and NICS have been calculated and analyzed. The analysis of the data showed that 3-chlorothiophene has the smallest values of the F_n coefficient. Therefore, it can be suggested that double bonds in 3-chlorothiophene are more delocalized. Electrochemical stability of 3-chlorothiophene and 3,4-dichlorothiophene is greater than other chlorothiophenes. Thermal stability and Zero-Point Energy of thiophene are greater than all chlorothiophenes. The 3-chlorothiophene and 3, 4-dichlorothiophene possesses the largest NICS value; these compounds have the largest electrical current in the thiophene ring. Accordingly, compound **B** (3-chlorothiophene) is the best candidate monomer among all chlorothiophenes for the synthesis of corresponding conducting polymers with modified characteristics. The synthesis feasibility, experimental measurement and thermo chemical stability of these chlorothiophenes are interesting subjects for study.

Acknowledgements

This research was supported by the Islamic Azad University, Toyserkan Branch, Toyserkan, Iran.

Author details

¹Young Researchers Club, Islamic Azad University, Toyserkan Branch, Toyserkan, Iran. ²Department of Chemistry, Islamic Azad University, Toyserkan Branch, Toyserkan, Iran.

Authors' contributions

SJ-B proposed and supervised the program, and participated in data analysis and draft revision, coordinated the study and modified the text. HS participated in data collection and manuscript drafting, completed the molecular design, the arrangement of the work and modification of the manuscript. SJ-B and HS participated in the separation and purification of compounds, acquisition of data, and collection of literature and helped to draft the manuscript. Authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 17 September 2010 Accepted: 14 March 2011
 Published: 14 March 2011

References

- Jameh-Bozorghi S, Shirani ILH: Theoretical study on the electronic, structural, properties and reactivity of a series of mono-, di-, tri- and tetrafluorothiophenes as well as corresponding radical cation forms. *J Fluor Chem* 2011.
- MacDiarmid AG: Synthetic metals: A novel role for organic polymers. *Chem Int E* 2001, **40**:2581-2585.
- Hall AW, Blackwood KM, Milne PE, Goodby JW: Novel UV cured coatings and adhesives based on the photoinitiated cyclopolymerization of derivatives of diallylamine. *Chem Commun* 2003, 2530-2531.
- Neves S, Gazotti WA, De Paoli M-A: Conducting Polymeric Nanomaterials. In *In Encyclopedia of Nanoscience and Nanotechnology. Volume 2*. Edited by: Nalwa HS. Los Angeles: American Scientific Publishers; 2004:133-152.
- Gangopadhyay R: Conducting Polymer Nanostructures. In *In Encyclopedia of Nanoscience and Nanotechnology. Volume 2*. Edited by: Nalwa HS. Los Angeles: American Scientific Publishers; 2004:105-131.
- Wallace GG, Innis PC, Kane-Maguire LAP: Inherently Conducting Polymer Nanostructures. In *In Encyclopedia of Nanoscience and Nanotechnology. Volume 4*. Edited by: Nalwa HS. Los Angeles: American Scientific Publishers; 2004:113-130.
- Epstein AJ: Introduction to Electronic Polymers: Influence of Nanostructure on Electronic Phenomena. In *In Organic Electronic Materials: Conjugated Polymers and Low Molecular Weight Organic Solids. Volume 41*. Edited by: Farchioni R, Grosso G. Amsterdam: Springer; 2001:3-39.
- Kolla HS, Surwade SP, Zhang X, MacDiarmid AG, Manohar SK: Absolute molecular Weight of Polyaniline. *J Am Chem Soc* 2005, **127**:16770-16771.
- Richard SP, Rajasekhar M, Subramania A: Synthesis of polythiophene nanoparticles by surfactant - assisted dilut polymerization method for high performance redox supercapacitors. *Int J Electrochem Sci* 2009, **4**:1289-1301.
- Bai H, Shi G: Gas sensors based on conducting polymers. *Sensors* 2007, **7**:267-307.
- Henry D, Tran YW, Julio MD, Richard BK: Toward an understanding of the formation of conducting polymer nanofibers. *ACS Nano* 2008, **2**:1841-1848.
- Ohkawa S, Ohta R, Kawabata K, Goto H: Polymerization in liquid crystal medium: preparation of polythiophene derivatives bearing a bulky pyrimidine substituent. *Polymers* 2010, **2**:393-406.
- Viinikanoja A, Lukkari J, Aaritalo T, Laiho T, Kankare J: Phosphonic acid derivatized polythiophene: A building block for metal phosphonate and polyelectrolyte multilayers. *Langmuir* 2003, **19**:2768-2775.
- Grevin B, Rannou P: Electrochemistry: Arrays of polymer nanowires. *Nature Materials* 2004, **3**:503-504.
- Nishinaga T, Yamazaki D, Tateno M, Iyoda M, Komatsu K: Synthesis and properties of octithiophene dication sterically segregated by annelation with bicycle [2.2]octene units. *Materials* 2010, **3**:2037-2052.
- Lipton-Duffin JA, Miwa JA, Kondratenko M, Cicoira FB, Sumpter G, Meunier V, Perepichka DF, Rosei F: Step-by-step growth of epitaxially aligned polythiophene by surface-confined reaction. *PNAS* 2010, **107**:11200-11204.
- Ramanathan K, Bangar M, Yun M, Chen W, Myung N, Mulchandani A: Bioaffinity sensing using biologically functionalized conducting-polymer nanowire. *J Am Chem Soc* 2005, **127**:496-497.
- Wayne E, Jones Jr, Chiguma J, Johnson E, Pachamuthu A, Santos D: Electrically and thermally conducting nanocomposites for electronic applications. *Materials* 2010, **3**:1478-1496.
- Wan MX: Conducting Polymer Nanotubes. In *In Encyclopedia of Nanoscience and Nanotechnology. Volume 2*. Edited by: Nalwa HS. Los Angeles, American Scientific Publishers; 2004:153-169.
- Maddalena F, Meijer EJ, Asadi K, Leeuw DM, Blom PWM: Doping kinetics of organic Semiconductors investigated by field-effect transistors. *Appl Phys Lett* 2010, **97**:0433021-0433023.
- Yang L, Qi Y, Yuan X, Shen J, Kim J: Direct synthesis, characterization and catalytic application of SBA-15 containing heteropolyacid. *Journal of Molecular Catalysis A: Chemical* 2005, **229**:199-205.
- Wallace GG, Spinks GM, Kane-Maguire LAP, Teasdale PR: Synthesis and Properties of Polythiophenes. In *Conductive electroactive polymers: intelligent polymer systems* Queensland: CRC Press; 2009, 197-226.
- Lukkari J, Viinikanoja A, Paukkunen J, Salomaki M, Mervi Janhonen, Aaritalo T, Kankare J: Oxidation induced variation in polyelectrolyte multilayers prepared from sulfonated self-dopable poly (alkoxythiophene). *Chem Commun* 2000, 571-572.

24. Asami R, Fuchigami T, Atobe M: **Development of a novel environmentally friendly electropolymerization of water-insoluble monomers in aqueous electrolytes using acoustic emulsification.** *Langmuir* 2006, **22**:0258-10263.
25. Chen J, David LO, Jennifer MP, Douglas RMF, Chee OT, Gordon GW: **Photoelectrochemical solar cells based on polyterthiophenes containing porphyrins using ionic liquid electrolyte.** *Electrochemical and Solid-State Letters* 2005, **8**:A528-A530.
26. Silvestri F, Marrocchi A: **Acetylene-Based materials in organic photovoltaics.** *Int J Mol Sci* 2010, **11**:1471-1508.
27. Oterom TF, Azelain EL: **Synthesis and properties of polythiophenes.** *Polym Comm* 1988, **29**:21-50.
28. Alvaro AA, Rosa LT: **Study of kinetic formation and the electrochemical behavior of polypyrrole films.** *J Chil Chem Soc* 2009, **54**:14-19.
29. Wallace GG, Spinks GM, Kane-Maguire LAP, Teasdale PR: *In Conductive electroactive polymers: intelligent polymer systems* Queensland: CRC Press; 2009, 77-107.
30. Frisch MJ, Trucks GW, Schlegel HB, Scuseria GE, Robb MA, Cheeseman JR, Zakrzewski VG, Montgomery JA, Startmann RE, Burant JC, Dapprich S, Millam JM, Daniels AD, Kudin KN, Strain MC, Farkas O, Tomasi J, Barone V, Cossi M, Cammi R, Mennucci B, Adamo C, Clifford S, Ochterski J, Petersson GA, Ayala PY, Morokuma QCK, Malick DK, Rabuck AD, Raghavachar K, et al: *GAUSSIAN 98 (Revision A3)*, Gaussian Inc Pittsburgh: PA; 1998.
31. Gross EK, Dreiz RM: *Density Functional Theory* New York: Plenum Press; 1995.
32. Carlier P, Deora N, Lam P, Harmon J, Williams L: *Carlier group gaussian user manual* 2010 [http://www.gaussian.com], Version 1.22.
33. Andersson MP, Uvdal P: **New scale factors for harmonic vibrational frequencies using the B3LYP density functional method with the triple-Basis Set 6-311+G(d,p).** *J Phys Chem A* 2005, **109**:2937-2941.
34. Omrani A, Sabzyan H: **Theoretical study of chloropyrroles as monomers for new conductive Polymers.** *J Phys Chem A* 2005, **109**:8874-8879.
35. Sabzyan H, Omrani A: **Ab initio and DFT study of all Mono-, Di-, Tri- and Tetrafluoropyrroles and their cations: predicting structural, spectroscopic, electropolymerization and electrochemical properties.** *J Phys Chem A* 2003, **107**:6476-6482.
36. Wallace GG, Spinks GM, Kane-Maguire LAP, Teasdale PR: **Processing and Device Fabrication.** *In Conductive electroactive polymers: intelligent polymer systems* Queensland: CRC Press; 2009, 231-245.
37. Nikoofard H, Sabzyan H: **DFT study of molecular structure and electronic properties of fluoromethylpyrrole oligomers including di-, tri- and tetramer.** *Journal of Fluorine Chemistry* 2007, **128**:668-673.
38. Kleinpeter E, Klod S: **Ab initio calculation of the anisotropy effect of multiple bonds and the ring current effect of arenes-application in conformational and configurational analysis.** *J Chem Soc Perkin Trans* 2001, **2**:1893-1898.
39. Martin NH, Caldwell BC, Carlson KP, Teague MR: **Ab initio calculation of through-space magnetic shielding of linear polycyclic aromatic hydrocarbons (acenes): Extent of aromaticity.** *J Mol Graph Model* 2009, **27**:689-692.
40. Chen Z, Wannere CS, Corminboeuf C, Puchta R, Schleyer PVR: **Nucleus-Independent Chemical Shifts (NICS) as an Aromaticity Criterion.** *Chem Rev* 2005, **105**:3842-3888.
41. Baldrige KK, Siegel JS: **Correlation of empirical (TMS) and absolute NMR chemical shifts predicted by ab initio computations.** *J Phys Chem A* 1999, **103**:4038-4042.
42. Zhang Y, Wu A, Xu Xin, Yan Y: **Geometric dependence of the B3LYP-Predicted magnetic shielding and chemical shifts.** *J Phys Chem A* 2007, **111**:9431-9437.
43. **Gaussian 03 Online Manual.** [http://www.nd.edu/~wschnei1/courses/CBE_547/g03_man/g_ur/k_nmr.htm].
44. Sabzyan H, Omrani A: **Ab initio and density functional study of electrical and thermochemical properties of mono-, di-, tri- and tetrafluoropyrroles and their cations and anions.** *J THEOCHEM* 2005, **713**:43-49.
45. Schleyer PVR, Manoharan M, Wang Z-X, Kiran B, Jiao H, Puchta R, Hommes NJRVE: **Dissected Nucleus-Independent Chemical Shift analysis of π -aromaticity and antiaromaticity.** *Org Lett* 2001, **3**:2465-2468.
46. Corminboeuf C, Heine T, Seifert G, Schleyer PVR, Weber J: **Induced magnetic fields in aromatic [n]-annulenes-interpretation of NICS tensor components.** *Phys Chem Chem Phys* 2004, **6**:273-276.
47. Alkorta I, Rozas I, Elguero J: **An ab initio study of the NMR properties (absolute shielding and NICS) of a series of significant aromatic and antiaromatic compounds.** *Tetrahedron* 2001, **57**:6043-6049.
48. Stanger A: **Nucleus-Independent Chemical Shifts (NICS): Distance Dependence and Revised Criteria for Aromaticity and Antiaromaticity.** *J Org Chem* 2006, **71**:883-893.
49. Fallah-Bagher-Shaidaei H, Wannere CS, Corminboeuf C, Puchta R, Schleyer PVR: **Which NICS aromaticity index for planar π -rings is best?** *Org Lett* 2006, **8**:863-866.

doi:10.1186/1752-153X-5-13

Cite this article as: Shirani IL Beigi and Jameh-Bozorghi: **Theoretical study on the electronic, structural, properties and reactivity of a series of mono-, di-, tri- and tetrachlorothiophenes as well as corresponding radical cation forms as monomers for conducting polymers.** *Chemistry Central Journal* 2011 **5**:13.

Publish with **ChemistryCentral** and every scientist can read your work free of charge

"Open access provides opportunities to our colleagues in other parts of the globe, by allowing anyone to view the content free of charge."

W. Jeffery Hurst, The Hershey Company.

- available free of charge to the entire scientific community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.chemistrycentral.com/manuscript/

