

RESEARCH ARTICLE

Open Access

Kinetics and mechanism of jack bean urease inhibition by Hg^{2+}

Nana Du¹, Mingming Chen^{1,2}, Zhaodi Liu¹, Liangquan Sheng^{1,2*}, Huajie Xu¹ and Shuisheng Chen¹

Abstract

Background: Jack bean urease (EC 3.5.1.5) is a metalloenzyme, which catalyzes the hydrolysis of urea to produce ammonia and carbon dioxide. The heavy metal ions are common inhibitors to control the rate of the enzymatic urea hydrolysis, which take the Hg^{2+} as the representative. Hg^{2+} affects the enzyme activity causing loss of the biological function of the enzyme, which threatens the survival of many microorganism and plants. However, inhibitory kinetics of urease by the low concentration Hg^{2+} has not been explored fully. In this study, the inhibitory effect of the low concentration Hg^{2+} on jack bean urease was investigated in order to elucidate the mechanism of Hg^{2+} inhibition.

Results: According to the kinetic parameters for the enzyme obtained from Lineweaver–Burk plot, it is shown that the K_m is equal to 4.6 ± 0.3 mM and V_m is equal to 29.8 ± 1.7 $\mu\text{mol NH}_3/\text{min mg}$. The results show that the inhibition of jack bean urease by Hg^{2+} at low concentration is a reversible reaction. Equilibrium constants have been determined for Hg^{2+} binding with the enzyme or the enzyme-substrate complexes ($K_i = 0.012$ μM). The results show that the Hg^{2+} is a noncompetitive inhibitor. In addition, the kinetics of enzyme inhibition by the low concentration Hg^{2+} has been studied using the kinetic method of the substrate reaction. The results suggest that the enzyme first reversibly and quickly binds Hg^{2+} and then undergoes a slow reversible course to inactivation. Furthermore, the rate constant of the forward reactions (k_{+0}) is much larger than the rate constant of the reverse reactions (k_{-0}). By combining with the fact that the enzyme activity is almost completely lost at high concentration, the enzyme is completely inactivated when the Hg^{2+} concentration is high enough.

Conclusions: These results suggest that Hg^{2+} has great impacts on the urease activity and the established inhibition kinetics model is suitable.

Keywords: Jack bean urease, Hg^{2+} ion, Inhibition, Kinetics

Background

Urease (urea amidohydrolase, EC 3.5.1.5) is a nickel-containing enzyme that catalyzes the hydrolysis of urea to ammonia and carbon dioxide. The enzyme is widely distributed in nature and is found in a variety of plants, algae, fungi, bacteria and soil [1-3]. Thus, urease plays an important role in the nitrogen metabolism of many microorganism and plants [4,5]. Despite the great importance of urease in biotechnology and medicine and also continuous interest in many laboratories [6-11], the mechanisms of hydrolysis of the metal active site of the

enzyme have not been studied in detail [12]. Study show that the cysteine is involved in the catalysis, as was demonstrated by reacting the enzyme with cysteine-reactive agents [13,14]. In view of the complexity of the role of the active site flap cysteine in the urease catalysis, effect of potential inhibitors on the reactivity of enzyme thiol groups, the active site flap thiol in particular, was investigated [15,16].

The main classes of urease inhibitors are: boron-containing compounds [7], thiol compounds [13], phosphate [15], phosphoroamide compounds [17], hydroxamic acids [14], F^- ions [18], and heavy metal ions [19,20]. Among the known inhibitors of urease, the inhibition of urease by heavy metal ions is said to result from the reaction of these ions with a sulfhydryl group in the active center of the enzyme by a reaction analogous to the formation of

* Correspondence: shenglq@fync.edu.cn

¹Fuyang Normal College, College of Chemistry and Chemical Engineering, Fuyang 236037, People's Republic of China

²Anhui University, College of Chemistry and Chemical Engineering, Hefei 230039, People's Republic of China

metal sulfides [21]. The formation of sulfides with the active center was confirmed experimentally by the correlation between the inhibitory efficiency of metal ions and the solubility products of their sulfides [22]. So, inhibition of urease by heavy metal ions can be studied to investigate the effect on the reactivity of enzyme thiol groups. Moreover, the inhibition of heavy metal ions is related to its biological toxicity, which causing loss of the biological function of the enzyme then affects the growth and survival of the animal and plants [20,22]. Studies published so far on inhibition of urease of both plant and bacterial origin by heavy metal ions have aimed either at investigating their toxicity or at detection of their trace amounts, e.g. of Hg^{2+} , Cu^{2+} and Ag^+ ions [23-27]. Mercury is a known biotoxin that can accumulate in the human body and show up in the food chain [28]. However, inhibitory kinetics of urease by low concentration Hg^{2+} has not been explored fully and only a few studies on the urease inhibition activities of heavy metal ions have appeared in the literature.

In this study, the inhibitory effect of the low concentration Hg^{2+} on jack bean urease was investigated. Our present investigation not only studied the kinetics of inhibition of the jack bean urease by the low concentration Hg^{2+} specifically but also researched related inhibition mechanisms. The results show that Hg^{2+} has great impacts on the jack bean urease activity. The inhibition of jack bean urease by the low concentration Hg^{2+} is shown to be reversible. In addition, the rate constants of inactivation are determined.

Results

Effect of Hg^{2+} concentration on urease activity

Urease (6 $\mu\text{g}/\text{ml}$, 0.02 M $\text{KH}_2\text{PO}_4\text{-K}_2\text{HPO}_4$) was incubated with different concentrations of Hg^{2+} for 15 min. Then, mixed solutions of urease and Hg^{2+} were added into the assay media (0.67 mM urea) to investigate the relation of Hg^{2+} concentration and the residual activity (Figure 1). In certain range, the activity of the enzyme decreases with increase of Hg^{2+} concentration. The IC_{50} value is estimated to be $0.018 \pm 0.001 \mu\text{M}$ when the Hg^{2+} concentration could lead to 50% of enzyme activity loss. The enzyme activity is almost completely lost when the Hg^{2+} concentration reach 0.08 μM .

The inhibition mechanism of urease by Hg^{2+}

The inhibition mechanism on the urease by Hg^{2+} was studied. The procedure used to obtain relationship between the enzyme activity and urease concentration was same as in the residual activity experiment. Figure 2 shows the relationship between the enzyme activity and urease concentration. The plots of the remaining enzyme activity against the enzyme concentrations in the presence of different Hg^{2+} concentrations give a family




Figure 1 Effect of Hg^{2+} on jack bean urease activity. Conditions were 1 mL assay system containing 0.02 M $\text{KH}_2\text{PO}_4\text{-K}_2\text{HPO}_4$ buffer (pH 7.4), 0.67 mM of urea, 6 $\mu\text{g ml}^{-1}$ of enzyme and different concentrations of Hg^{2+} at 25°C for 15 min.

of straight lines, which all pass through the origin, indicating that the inhibition of Hg^{2+} on the enzyme is reversible [29]. Increasing the inhibitor concentration resulted in a descending slope of the line. The presence of inhibitor does not bring down the amount of effective enzyme, but just result in the inhibition and decreasing of activity of the enzyme.

In this section, the Hg^{2+} concentration was held at different constant values (lower than for complete inactivation), while the substrate concentrations were varied, permitting measurement of the effect of increasing substrate concentration [S] on the initial reaction rate (ν). Plots of $1/\nu$ vs. $1/[S]$ for different concentrations of inhibitor (Figure 3) show that Hg^{2+} is a noncompetitive





Figure 2 The relationship between the enzyme activity and urease concentration. The assay conditions were as described for Figure 1, except that the enzyme concentration. Numbers denote Hg^{2+} concentration (μM).




inhibitor [30]. Increasing the Hg^{2+} concentration results in a family of lines with a common intercept on the $1/[S]$ axis but with different slopes and intercepts on the $1/v$ axis, indicating that K_m is unchanged by the presence of different inhibitor concentrations while V_m is changed. The equilibrium constant for the inhibitor binding, K_i , can be obtained from the plot of $1/V_m$ versus the Hg^{2+} concentration, as shown in the inset. The inhibition constant, K_i , for Hg^{2+} obtained from the experimental data is $0.012 \mu M$.

Determination of the kinetic parameters of urease

The kinetic behavior of jack bean urease in the hydrolysis of urea has been studied specifically. Under the conditions employed in the present study, the hydrolysis of urea by urease follows Michaelis-Menten kinetics. The kinetic parameters for the enzyme have been obtained from Lineweaver-Burk plot (Figure 3, curve 0) with K_m ($4.6 \pm 0.3 \text{ mM}$) and V_m ($29.8 \pm 1.7 \mu\text{mol NH}_3/\text{min mg}$) which are in agreement with previously obtained values [31,32].

Kinetics of the substrate reaction in the presence of different concentrations of Hg^{2+}

Figure 4(a) shows the time process of the hydrolysis of the substrate without enzyme-inhibitor preincubation. The assay mixture (1.0 ml) contained 40 mg L^{-1} urea, $6 \mu\text{g ml}^{-1}$ urease and different concentrations Hg^{2+} in $0.02 \text{ M KH}_2\text{PO}_4\text{-K}_2\text{HPO}_4$ buffer ($\text{pH}=7.4$). At each Hg^{2+} concentration, the rate decreases with increasing time until a straight line is approached, is accord with the conclusion in ref.22. The results analyzed by Tsou/s method [33] suggest that the formation of the enzyme Hg^{2+} complex is a reversible reaction at low Hg^{2+} concentration [29], while the formation of the enzyme Hg^{2+} complex is irreversible reaction at $1.0 \mu\text{M}$ Hg^{2+} because the asymptote becomes to parallel to abscissa without residual enzyme activity [22]. In this paper, our chief aim is to investigate the reversible inhibition kinetics of the enzyme by Hg^{2+} at low concentration. According to Eq. (6), plots of $\ln([P]_{\text{calc}} - [P]_t)$ versus t give a family of straight lines at different concentrations of Hg^{2+} with slopes of $-A$



(Figure 4(b)), the apparent rate constants of inhibition. The microscopic rate constant of the reverse inactivation of the enzyme (k_{-0}) in these experimental conditions are listed in Table 1. The value of k_{-0} is almost the same for different Hg^{2+} concentrations. From Eq. (7), a plot of $1/(A-k_{-0})$ versus $1/[\text{Hg}]$ gave a straight line, which is fit using a least squares analysis. The straight line with a slope of K_i/k_{+0} intercepts on the ordinate with $1/k_{+0}$. The microscopic rate constant, k_{+0} , and the equilibrium constant for inhibitor binding, K_i , obtained from these values is $0.013 \mu\text{M}$.

Kinetics of the reaction at different substrate concentrations in the presence of Hg^{2+}

Figure 5(a) shows the kinetic courses of substrate reaction at different urea concentrations in the presence of $0.005 \mu\text{M} \text{Hg}^{2+}$. At each substrate concentration, the rate decreases with increasing time until a straight line is approached. The initial rate and the slope of the asymptote increase with increasing substrate concentration. Similarly, plots of $\ln([P]_{\text{calc}} - [P]_t)$ against t give a family of straight lines at different concentrations of the substrate with slopes of $-A$ (Figure 5(b)). It can be obtained through suitable plots for the apparent forward rate constants, A . A plot of the slopes of the straight lines in Figure 5(b) versus substrate concentration $[S]$ gives a horizontal straight line, Figure 5(c), indicating that the substrate concentration $[S]$ does not affect the microscopic rate constants: k_{+0} and k_{-0} . The results are shown in Table 2.

Discussion

In this paper, the effects of low concentration Hg^{2+} on the enzyme activity were studied specifically, the results of which show that the inhibitor concentration leading to 50% of enzyme activity loss, IC_{50} , is estimated to be $0.018 \pm 0.001 \mu\text{M}$. In addition, kinetics of inhibition of the enzyme by Hg^{2+} at the low concentration was also studied by the substrate reaction kinetic method described by Tsou [33]. As a result, the kinetic parameters for the enzyme obtained from Lineweaver–Burk plot show that K_m is equal to $4.6 \pm 0.3 \text{ mM}$ and V_m is equal to 29.8 ± 1.7

Table 1 Microscopic inhibitory rate constants of the jack bean urease in different concentrations [S]

[S] (mg L ⁻¹)	k_{+0} (min ⁻¹)	k_{-0} (min ⁻¹)
20		0.036
30		0.037
40	0.267^I 0.303^{II}	0.036
50		0.035

^I From the intercept of the straight line of $1/(A-k_{-0})$ vs. $1/[\text{Hg}]$. ^{II} From the slope of the straight line of $1/(A-k_{-0})$ vs. $1/[\text{Hg}]$.




Figure 5 Inhibition kinetics for jack bean urease in various concentrations of substrate. (a) Course of the substrate reaction. Assay conditions were as described for Figure 1. Numbers denote urea concentration (mg L⁻¹). (b) Semilogarithmic plot of $\ln([P]_{\text{calc}} - [P]_t)$ vs. t with the data taken from curves in (a). (c) The apparent forward inactivation rate constants (A) vs. urea concentrations in the presence of Hg^{2+} .

$\mu\text{mol NH}_3/\text{min mg}$, which generally accord with the values reported in literature [31,32]. When the reaction time is sufficiently long, the concentration of product

Table 2 Microscopic rate constants of jack bean urease by different concentrations Hg²⁺

[Hg] (μM)	A(min ⁻¹)	k ₊₀ (min ⁻¹)	k ₋₀ (min ⁻¹)
0.0050	0.118	0.279	0.036
0.0125	0.173	0.267	0.037
0.0200	0.228	0.307	0.036

approaches a constant final value, which decreases as the concentration of Hg²⁺ increases and is accord with the conclusion in ref.22. The treated enzyme still has partial residual activity at low concentration of Hg²⁺, which is analyzed by the method described by Tsou [33] and also suggests that inactivation of urease by the low concentration Hg²⁺ is a reversible reaction under the experimental conditions [29], while the formation of the enzyme Hg²⁺ complex is an irreversible reaction at 1.0 μM Hg²⁺ because the asymptote tends to parallel to the abscissa without residual enzyme activity [22]. The plots of the remaining enzyme activity against the enzyme concentrations in the presence of Hg²⁺ with different concentrations give a family of straight lines, all of which pass through the origin, indicating that the inhibition of Hg²⁺ on the enzyme is reversible [29]. The results suggest that the enzyme first reversibly and quickly binds Hg²⁺ and then undergoes a slow reversible course to inactivation on the basis of the previous study [22]. Plots of 1/v against 1/[S] at different concentrations of Hg²⁺ give a family straight line with a common intercept on the 1/[S] axis but with different slopes and intercepts on the 1/v axis, which indicates that Hg²⁺ is a noncompetitive inhibitor for the enzyme [30].

In accordance with the microscopic rate constants are determined and shown in Table 1 and Table 2, we can see that the values of k₊₀ are almost the same (0.279, 0.267, 0.307 min⁻¹) and the values of k₋₀ are the same (0.036, 0.037, 0.036 min⁻¹) as well. Furthermore, the microscopic rate constants of the forward and the reverse reactions slightly vary with the increasing concentration of the substrate. The results indicate that the presence of substrate does not provide protection for the enzyme against the inhibition by Hg²⁺. Besides, the rate constant for forward inactivation (k₊₀) is much larger than that for reverse reactivation (k₋₀). According to combination with the fact that the enzyme activity is almost completely lost when the concentration of Hg²⁺ reaches 0.08 μM, the enzyme is completely inactivated when the concentration of Hg²⁺ is high enough. The results suggest that Hg²⁺ has great impacts on the urease activity. Based on the fact that thiols are involved in binding Hg that was provided in the previous study, there is another demonstration that sulfhydryl group is essential for the activity of the enzyme.

Experimental

Materials


Jack bean urease, Fluka, with specific activity 57 μmol NH₃/min mg protein was used (without further purified). Urea (substrate), mercuric chloride (inhibitor) and other reagents are local products of analytical grade. The water used was redistilled and ion-free.

Assay

The standard residual activity assay mixture (1.0 ml) contained 0.67 mM urea in 0.02 M KH₂PO₄-K₂HPO₄ buffer (pH 7.4). Urease was incubated by Hg²⁺ for 15 min. Reactions were initiated by the addition of small aliquots of the enzyme and Hg²⁺ mixed solution, typically 3 μl containing 2 mg ml⁻¹ enzyme. After reaction for 15 at 25°C, 2 ml phenol solution and 3 ml hypochlorite sodium solution were added into the mixture at 35°C.

Progress curves of the enzyme reactions were obtained by measuring the ammonia generated as a function of reaction time. The assay mixture (1.0 ml) typically contained 20-50 mg L⁻¹ urea in 0.02 M KH₂PO₄-K₂HPO₄ buffer (pH 7.4), 0.005-0.025 μM Hg²⁺ and 6 μg ml⁻¹ urease. Reactions were initiated by adding urease to the mixture of urea and Hg²⁺. After reaction for some time at 25°C, 2 ml phenol solution and 3 ml hypochlorite sodium solution were added into the mixture at 35°C.

The reaction was monitored by measuring the ammonia concentration by the phenol-hypochlorite method [34] in samples, which were removed from the reaction mixtures at time intervals. Absorption at 625 nm was recorded using a spectrophotometer. One unit (U) of enzymatic activity was defined as the amount of enzyme required to produce 1μM ammonia per min under these



conditions. The activity of uninhibited urease was accounted as the control activity of 100%.

Determination of microscopic rate constants

The progress-of-substrate-reaction method that previously described by Tsou [33] was used to study the inhibitory kinetics of jack bean urease by Hg^{2+} . The substrate reaction progress curve was analyzed to obtain the reaction rate constants as detailed below. The reaction was carried out at a constant temperature of 25°C.

For slow, reversible inhibition with fractional residue activity, the kinetic model of the enzyme reacting with the substrate and the inhibitor can be written as Scheme 1 [30], where S, P, Hg and E respectively denote substrate, product, inhibitor (Hg^{2+}) and enzyme. EHg, ES and ESHg are the respective complexes. E'Hg and E'HgS are inactive enzyme forms. K_i is the inhibition constant for the inhibitor (Hg^{2+}). k_{+0} and k_{-0} are rate constants for forward and reverse inactivation of the enzyme, respectively.

As is usual the case $[\text{S}] \gg [\text{E}_0]$ and that the modification reactions are relatively slow compared with the set up of the steady-state of the enzymatic reaction. The product formation can be written as:

$$[P]_t = \frac{\nu k_{-0}}{A} t + \frac{(A - k_{-0})\nu}{A^2} (1 - e^{-At}) \quad (1)$$

$$A = \frac{k_{+0}[\text{Hg}]}{K_i + [\text{Hg}]} + k_{-0} \quad (2)$$

$$B = k_{-0} \quad (3)$$

where $[P]_t$ is the concentration of the product formed at time t , which is the reaction time. A and B are the apparent rate constants for inactivation and reactivation, respectively. k_{+0} and k_{-0} are the microscopic rate constants for the forward and reverse reactions, $[\text{S}]$ and $[\text{Hg}]$ are the concentrations of the substrate and inhibitor. ν is the initial rate of reaction in the absence of the inhibitor, where $\nu = V_m \times [\text{S}] / (K_m + [\text{S}])$, which is the Michaelis-Menten equation. When t is sufficiently long, the curves become straight lines and the product concentration is written as $[P]_{\text{calc}}$:

$$[P]_{\text{calc}} = \frac{\nu k_{-0}}{A} t + \frac{(A - k_{-0})\nu}{A^2} \quad (4)$$

A plot of $[P]_{\text{calc}}$ vs. t gives a straight line. Combining Eqs. (1) and (4) gives:

$$[P]_{\text{calc}} - [P]_t = \frac{(A - k_{-0})\nu}{A^2} e^{-At} \quad (5)$$

$$\ln([P]_{\text{calc}} - [P]_t) = \ln[\nu(A - k_{-0})/A^2] - At \quad (6)$$

where $[P]_{\text{calc}}$ is the product concentration to be expected from the straight-line portions of the curves as

calculated from Eq. (4) and $[P]_t$ is the product concentration actually observed at time t . Plots of $\ln([P]_{\text{calc}} - [P]_t)$ versus t give a series of straight lines at different concentrations of inhibitor ($[\text{Hg}]$) with slopes of $-A$. The apparent forward rate constant A can be obtained from such graphs. From Eq. (4), a plot of $[P]_{\text{calc}}$ against time, t , gives a straight line with a slope of $\nu k_{-0}/A$. From the slope of the straight line, k_{-0} can be obtained.

The apparent forward rate constant, A , is independent of the substrate concentration, but depends on the inhibitor concentration From Eq. (2) and Eq. (3):

$$\frac{1}{A - k_{-0}} = \frac{K_i}{k_{+0}[\text{Hg}]} + \frac{1}{k_{+0}} \quad (7)$$

Plot of $1/(A - k_{-0})$ versus $1/[\text{Hg}]$ gives a straight line with a slope of K_i/k_{+0} and an intercept of $1/k_{+0}$ on the ordinate and $-1/K_i$ on the abscissa which can be used to determine the microscopic rate constant, k_{+0} , and the equilibrium constant for the inhibitor binding, K_i .

Conclusions

The inhibitory effect of Hg^{2+} on jack bean urease was investigated that the Hg^{2+} displayed strong inhibitory activity against jack bean urease. The experiment result shows that the inhibition of urease by low concentration Hg^{2+} is a reversible reaction and the inhibition belongs to be noncompetitive. The microscopic rate constants of the forward and the reverse reactions vary little with increasing substrate concentration, indicating that presence of substrate does not offer protection of this enzyme against inhibition by Hg^{2+} . The data fit well to model, which indicates that the established inhibition kinetics model is suitable.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

ND made a significant contribution to acquisition of data, analysis and manuscript preparation. MC has made a substantial contribution to experimental design and data analysis. ZL and HX participated in partial experiments. LS made a significant contribution to experimental design, data analysis, and manuscript revision. SC participated in study design and manuscript revision. All authors read and approved the final manuscript.

Acknowledgement

This work was financially supported by the Nature Science Foundation of China (20971024), the International Sea Area Resources Survey and Development of the 12th Five-year Plan of China (DY125-15-E-01) and the Higher Education Institutions Key Nature Science Foundation of Anhui (kj2009A127).

Received: 11 October 2012 Accepted: 30 November 2012

Published: 10 December 2012

References

1. Karplus PA, Pearson MA, Hausinger RP: 70 Years of Crystalline Urease: What Have We Learned? *Acc Chem Res* 1997, **30**:330-337.
2. Krajewska B: Ureases I. Functional, catalytic and kinetic properties: A review. *J Mol Catal B: Enzym* 2009, **59**:9-21.

- Das N, Kayastha AM, Srivastava PK: Purification and characterization of urease from dehusked pigeonpea (*Cajanus cajan* L.) seeds. *Phytochem* 2002, **61**:513–521.
- Dong XW, Li YG, Li ZW, Cui YM, Zhu HL: Synthesis, structures and urease inhibition studies of copper(II) and nickel(II) complexes with bidentate N, O-donor Schiff base ligands. *J Inorg Biochem* 2012, **108**:22–29.
- Murafuji T, Azuma T, Miyoshi Y, Ishibashi M, Mustafizur Rahman AFM, Migita K, Sugihara Y, Mikata Y: Inhibition of jack bean urease by organobismuth compounds. *Bioorg Med Chem Lett* 2006, **16**:1510–1513.
- You ZL, Shi DH, Zhang JC, Ma Y, Wang C, Li K: Synthesis, structures, and urease inhibitory activities of oxovanadium(V) complexes with Schiff bases. *Inorg Chim Acta* 2012, **384**:54–61.
- Benini S, Rypniewski WR, Wilson KS, Mangani S, Ciurli S: Molecular details of urease inhibition by boric acid: insights into the catalytic mechanism. *J Am Chem Soc* 2004, **126**:3714–3715.
- Mao WJ, Lv PC, Shi L, Li HQ, Zhu HL: Synthesis, molecular docking and biological evaluation of metronidazole derivatives as potent *Helicobacter pylori* urease inhibitors. *Bioorg Med Chem* 2009, **17**:7531–7536.
- You ZL, Lu Y, Zhang N, Ding BW, Sun H, Hou P, Wang C: Preparation and structural characterization of hetero-dinuclear Schiff base copper(II)-zinc (II) complexes and their inhibition studies on *Helicobacter pylori* urease. *Polyhedron* 2011, **30**:2186–2194.
- Mari AJ, Domi N, Carmen SN, Mari AF, Juan AP, Sara SF, Oscar U, Fabrice H, Jose M, Garci AM: Design, synthesis, and biological evaluation of phosphoramidate derivatives as urease inhibitors. *J Agric Food Chem* 2008, **56**:3721–3731.
- Cheng K, Zheng QZ, Zhu HL: Syntheses, structures and urease inhibitory activities of mononuclear cobalt(II) and 1D cobalt(II) complexes with ligands derived from 3-formylsalicylic acid. *Inorg Chem Commun* 2009, **12**:1116–1119.
- Krajewska B, Eldik RV, Brindell M: Temperature- and pressure-dependent stopped-flow kinetic studies of jack-bean urease. Implications for the catalytic mechanism. *J Biol Inorg Chem* 2012, **17**:1123–1134.
- Krajewska B, Zaborska W: Double mode of inhibition-inducing interactions of 1,4-naphthoquinone with urease: arylation versus oxidation of enzyme thiols. *Bioorgan Med Chem* 2007, **15**:4144–4151.
- Krajewska B, Zaborska W: Jack bean urease: The effect of active-site binding inhibitors on the reactivity of enzyme thiol groups. *Bioorg Chem* 2007, **35**:355–365.
- Zaborska W, Krajewska B, Kot M, Karcz W: Quinone-induced inhibition of urease: Elucidation of its mechanisms by probing thiol groups of the enzyme. *Bioorg Chem* 2007, **35**:233–242.
- Krajewska B, Ciurli S: Jack bean (*Canavalia ensiformis*) urease. Probing acid-base groups of the active site by pH variation. *Plant Physiol and Bioch* 2005, **43**:651–658.
- Benini S, Rypniewski WR, Wilson KS, Miletti S, Ciurli S, Mangani S: A new proposal for urease mechanism based on the crystal structures of the native and inhibited enzyme from *Bacillus pasteurii*: Why urea hydrolysis costs two nickels. *Structure* 1999, **7**:205–216.
- Tarun EI, Rubinov DB, Metelitzta DI: Inhibition of urease by cyclic b-triketones and fluoride ions. *Appl Biochem Micro* 2004, **40**:337–344.
- Behbehani GR, Saboury AA, Taherkhani A, Barzegar L, Mollaagazade A: A thermodynamic study on the binding of mercury and silver ions to urease. *J Therm Anal Calorim* 2011, **105**:1081–1085.
- Krajewska B: Mono-(Ag, Hg) and di-(Cu, Hg) valent metal ions effects on the activity of jack bean urease, Probing the modes of metal binding to the enzyme. *J Enzyme Inhib Med Chem* 2008, **23**:535–542.
- Takishima K, Suga T, Mamiya G: The structure of jack bean urease. The complete amino acid sequence, limited proteolysis and reactive cysteine residues. *Eur J Biochem* 1988, **175**:151–165.
- Krajewska B, Zaborska W, Chudy M: Multi-step analysis of Hg²⁺ ion inhibition of jack bean urease. *J Inorg Biochem* 2004, **98**:1160–1168.
- Chen JL, Zheng AF, Chen AH, Gao YC, He CY, Kai XM, Wu GH, Chen Y: A functionalized gold nanoparticles and rhodamine 6G based fluorescent sensor for high sensitive and selective detection of mercury(II) in environmental water samples. *Anal Chim Acta* 2007, **599**:134–142.
- Shi R, Stein K, Schwedt G: Determination of mercury(II) traces in drinking water by inhibition of an urease reactor in a flow injection analysis (FIA) system. *Fresenius J Anal Chem* 1997, **357**:752–755.
- Preininger C: The enzymatic determination of mercury and copper using acid urease, The effects of buffers. *Mikrochim Acta* 1999, **130**:209–214.
- Prakasha O, Talat M, Hasan SH: Response surface design for the optimization of enzymatic detection of mercury in aqueous solution using immobilized urease from vegetable waste. *J Mol Catal B Enzym* 2009, **56**:265–271.
- Pal P, Bhattacharyay D, Mukhopadhyay A, Sarkar P: The detection of mercury, cadmium, and arsenic by the deactivation of urease on rhodized carbon. *Environ Eng Sci* 2009, **26**:25–32.
- Lin JC, Xie XL, Gong M, Wang Q, Chen QX: Effects of mercuric ion on the conformation and activity of *Penaeus Vannamei* β-N-acetyl-d-glucosaminidase. *Int J Biol Macromol* 2005, **36**:327–330.
- Huang XH, Chen QX, Wang Q, Song KK, Wang J, Sha L, Guan X: Inhibition of the activity of mushroom tyrosinase by alkylbenzoic acids. *Food Chem* 2006, **94**:1–6.
- Zhan RQ, Chen QX, Zhen WZ, Lin JY, Zhuang ZL, Zhou HM: Inhibition kinetics of green crab (*Scylla serrata*) alkaline phosphatase activity by dithiothreitol or 2-mercaptoethanol. *Int J Biochem Cell B* 2000, **32**:865–872.
- Krajewska B, Zaborska W: The effect of phosphate buffer in the range of pH 5.80–8.07 on jack bean urease activity. *J Mol Catal B Enzym* 1999, **6**:75–81.
- Zaborska W, Krajewska B, Leszko M, Olech Z: Inhibition of urease by Ni²⁺ ions Analysis of reaction progress curves. *J Mol Catal B Enzym* 2001, **13**:103–108.
- Tsou CL: Kinetics of substrate reaction during irreversible modification of enzyme activity. *Adv Enzymol Relat Areas Mol Biol* 1988, **61**:381–436.
- Sequeira M, Diamond D, Daridon A, Lichtenberg J, Verpoorte S, Rooij NF: Progress in the realisation of an autonomous environmental monitoring device for ammonia. *Trends in Anal Chem* 2002, **21**:816–827.

doi:10.1186/1752-153X-6-154

Cite this article as: Du et al.: Kinetics and mechanism of jack bean urease inhibition by Hg²⁺. *Chemistry Central Journal* 2012 **6**:154.

Publish with **ChemistryCentral** and every scientist can read your work free of charge

“Open access provides opportunities to our colleagues in other parts of the globe, by allowing anyone to view the content free of charge.”

W. Jeffery Hurst, The Hershey Company.

- available free of charge to the entire scientific community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
<http://www.chemistrycentral.com/manuscript/>



ChemistryCentral