

RESEARCH ARTICLE

Open Access

Deep investigation on inorganic fraction of atmospheric PM in Mediterranean area by neutron and photon activation analysis

Pasquale Avino^{1*}, Geraldo Capannesi², Maurizio Manigrasso¹, Alberto Rosada² and Mario Vincenzo Russo³

Abstract

Background: Anthropogenic activities introduce materials increasing levels of many dangerous substances for the environmental quality and being hazardous to human health. Major attention has been given to those elements able to alter the environment and endanger human health.

The airborne particulate matter pollutant is considered one of the most difficult task in environmental chemistry for its complex composition and implications complicating notably the behavior comprehension. So, for investigating deeply the elemental composition we used two nuclear techniques, Neutron Activation Analysis and Photon Activation Analysis, characterized by high sensitivity, precision and accuracy. An important task has been devoted to the investigation of Quality Control (QC) and Quality Assurance (QA) of the methodology used in this study.

This study was therefore extended as far back as possible in time (from 1965 until 2000) in order to analyze the trend of airborne concentration of pollutant elements in connection with the industrial and lifestyle growth during the entire period.

Results: Almost all the elements may be attributed to long-range transport phenomena from other natural and/or anthropogenic sources: this behavior is common to all the periods studied even if a very light decreasing trend can be evidenced from 1970 to 2002. Finally, in order to investigate a retrospective study of elements in PM₁₀ and their evolution in relationship with the natural or anthropogenic origins, we have investigated the Enrichment Factors. The study shows the EF trends for some elements in PM₁₀ during four decades.

Conclusions: The two nuclear techniques have allowed to reach elevated sensibility/accuracy levels for determining elements at very low concentrations (trace and ultra-trace levels). The element concentrations determined in this study do not basically show a significant level of attention from a toxicological point of view.

Keywords: Element, PM₁₀, PM_{2.5}, Atmosphere, Urban air, Enrichment factor, INAA, IPAA

Background

One of the most interesting and difficult task in environmental chemistry is the investigation of the inorganic chemical composition of the particulate matter. Different reasons are the basis of this consideration: for instance, the very low levels of some elements (e.g., Cd, Hg, Se, at ultra-trace levels), the sampling collection which is more representative as possible, the availability of analytical methods (especially, no-destructive but really sensitive methodologies) [1-11].

The element distribution in air is fundamentally determined by resuspension from soil and water of various substances of natural and/or artificial origin, by their type of circulation due to the meteorological events and by the chemical element behavior. Anthropogenic activities introduce species increasing levels of many substances which may endanger the environmental quality and represent a hazard to human health. Major attention has been given to those elements able to alter the environment and endanger human health.

The issue regarding the determination of trace elements in airborne particulate have a preminent position due to the presence of some toxic elements (e.g., Cd, Hg and Pb): further, the airborne particulate matter enters

* Correspondence: p.avino@inail.it

¹DIPIA, INAIL settore Ricerca, Certificazione e Verifica, Via IV Novembre 144, Rome 00187, Italy

Full list of author information is available at the end of the article

into the climate global change problem, giving place to increasing global levels that may affect widely the biological systems.

The very complex composition and implications of the airborne particulate matter pollutant PM_{10} and, especially, the distribution and multielemental composition of the fine fraction (i.e., particles with diameter $<2.5 \mu\text{m}$, $PM_{2.5}$), complicate notably the behavior comprehension. The evaluation of background levels due to natural pathways of circulation, seems the preliminar action to be undertaken. This study was therefore extended as far back as possible in time (from the seventies until nowadays) in order to analyze the trend of airborne concentration of pollutant elements in connection with the industrial and lifestyle growth during the entire period.

Instrumental nuclear techniques are widely used in this field [12] as they represent the most reliable method for analyzing trace and/or ultra-trace elements in air particulate PM_{10} and $PM_{2.5}$. Instrumental Neutron Activation Analysis (INAA) as well Instrumental Photon Activation Analysis (IPAA) have been employed in this work to measure interesting toxicologically elements. INAA is not universally applicable with regard to the elements that can be determined, as the technique does not have sufficient sensitivity for Ca, Ti, Sr and Zr for example, and Nb, Pb, Tl and Y are impossible to measure. Similarly, determination of some elements is complicated due to interfering nuclear reactions or from the products of uranium fission (samples with elevated U concentration) [13,14]. An useful complementary method to INAA is Instrumental Photon Activation Analysis (IPAA), which enables the determination of the above elements. The IPAA method has been utilised on a much smaller scale in analytical field [15,16]. In IPAA, the nuclei can be activated through photonuclear process: in fact, contrary to INAA based mainly on the neutron capture reaction (n, γ), PAA employs photonuclear reactions, particularly the (γ, n) reaction [17]. Activation is induced by high energy photons having energy of at least 10 MeV. The photonuclear reaction data of the elements require around 30 MeV for activation energy. For analytical data interpretation, IPAA results have been considered only for the elements that cannot be determined and/or are difficult to determine by INAA [18-27], and for which IPAA provides results of comparable or better quality. Determination of other elements, such as Mn, Rb, Cs, Ba, Ce, U by IPAA should be considered of subsidiary value because INAA is more sensitive. Finally, as reported in literature [28], IPAA is not as commonly used as INAA but possibilities for element determination by both INAA and IPAA depend strongly on the matrix composition.

In particular, over As, Cd, Cr, Hg, Pb, Sb and Zn, i.e. metals considered of greater health concern, other elements, e.g. Au, Cs, La, Mo, Sc, Se, Sm, Th, W, were measured.

Results and discussion

Quality control

Table 1 shows the analytical quality control performed on a Standard Reference Material (SRM). This control was performed through an intercomparison campaign for 14 elements promoted by the International Atomic Energy Agency (IAEA) on air filter samples among different worldwide laboratories (130) using both spectrochemical (colorimetry, fluorescence, x-ray fluorescence, infrared spectrometry, atomic absorption and emission spectrometry, ICP-AES, ICP-MS), electrochemical (polarography, voltammetry) and nuclear (INAA, IPAA, isotopic dilution, beta counting) analytical techniques. For each element our values ("measured value") and the "certified value" are reported: the third column ("average value") represents the value averaged among all the determinations performed by different laboratories interested in the round-robin. As it can be noted, the agreement between our and the real value is quite good except for some elements such as Ba, U and Zn. For Ba and U this discrepancies can be due to the difficulty to analyze such kind of elements even if for Ba our "measured value" (43.4 ± 0.5) and the "average value" (39.05 ± 15.62) are quite similar. For Zn the situation is little bit different. The "measured value" ($132 \pm 18 \mu\text{g g}^{-1}$) falls out the certified value ($152 \mu\text{g g}^{-1}$) whereas it is in "average value" range ($141 \pm 16 \mu\text{g g}^{-1}$): the two-tailed P value is less than 0.001 and, by conventional criteria, this difference is considered to be statistically significant.

Table 1 Results of the Quality Control on IAEA air filter samples ($\mu\text{g g}^{-1}$)

Element	Measured value	Certified value	Average value	
	mean \pm s.d.	mean	mean	2σ (%)
As	4.9 ± 0.5	5.6	4.59	43
Au	1.26 ± 0.10	1.15	1.06	21
Ba	43.4 ± 0.5	53.8	39.05	40
Cd	10.6 ± 1.0	9.96	9.8	18
Co	1.3 ± 0.1	1.12	1.18	38
Cr	4.7 ± 0.8	5.6	4.8	13
Cu	51.6 ± 0.5	48.8	44.8	16
Fe	193 ± 17	207.9	200.1	8
Mn	31.2 ± 1.0	31.9	30.1	14
Mo	1.26 ± 0.2	1.14	1.56	70
Se	1.01 ± 0.10	1.06	1.01	11
U	0.78 ± 0.10	1.02	0.99	14
V	8.04 ± 0.35	8.00	7.2	16
Zn	132 ± 18	152	141	12

The "measured value" is the average of seven determinations on seven different replicates. s.d.: standard deviation.

Particulate matter results

Average concentration values, minimum and maximum levels and standard deviation of the elements determined in the PM_{2.5} fraction are shown in Table 2 whereas the correlation coefficients of the analyzed elements are reported in Table 3. It should be noted that many elements cannot be possible to determine in these samples: the main reasons depend on both the granulometric size fraction, 2.5 μm, as just reported in literature [9], and the very low levels reached by some elements (e.g., Nd is below LOD). Basically, the concentration levels of the elements are very low. There is only a very significant correlation between Br and Sb (correlation coefficient 0.95). The Br presence in air is essentially attributed to natural (e.g., marine aerosol) and anthropogenic (e.g., automotive traffic [7]) sources. This high correlation coefficient with Sb hypothesizes a strong anthropogenic contribution to the Br level. Furthermore, it

Table 2 Synoptic table (mean value, min-max values and standard deviation) of elements concentration (ng m⁻³) determined in PM_{2.5} in downtown Rome (LOD: limit of detection; * expressed as pg m⁻³; ** expressed as μg m⁻³)

Element	PM2.5		
	Mean	Min-Max	St. dev.
As	1.06	0.121-2.76	0.044
Au	0.009	0.000-0.050	0.012
Ba	3.76	1.91-6.45	2.38
Br	17.1	3.20-50.4	13.9
Ce	0.130	0.033-0.335	0.089
Co	0.167	0.077-0.331	0.065
Cr	3.03	1.29-6.40	1.30
Cs	0.047	0.004-0.124	0.037
Eu*	1.14	1.12-1.16	0.029
Fe**	0.074	0.005-0.212	0.059
Hf	0.018	0.006-0.032	0.010
Hg	0.818	0.195-2.12	0.655
La*	22.6	8.73-53.3	10.5
Mo	0.748	0.017-3.04	0.699
Nd	<LOD		
Ni	3.54	1.91-5.82	1.45
Rb	1.82	0.416-3.74	1.07
Sb	3.60	0.690-12.6	3.24
Sc*	3.14	0.208-7.49	2.41
Se	0.567	0.116-1.55	0.415
Sm*	3.88	0.208-7.78	2.13
Th	0.027	0.007-0.041	0.010
W	0.636	0.062-2.86	0.682
Yb	0.011	0.003-0.027	0.007
Zn	58.0	4.78-252	61.3

can be noted a wide scattering between the correlation coefficients: only As, Co, Fe, Sc, Sb, Se are highly correlated ($0.7 < \text{correlation coefficient} < 0.9$, marked in bold in Table 3) whereas Au, Ba, Br, Ce, Cr, Cs, La, Sm result scarcely correlated. This behavior can be expected considering the chemical-physical properties of the elements and the granulometric size (<2.5 μm), as previously evidenced in previous papers [7,9].

In Tables 4 and 5 are reported for each element both the summer and winter average levels as well as the maximum and minimum values for each season measured in PM₁₀ samples by INAA and IPAA. The samples were collected in outskirts of Rome and downtown [29], respectively, during summer and winter period. It may be noted that elements of anthropogenic origin show winter concentration levels higher than the summer ones, probably owing to an enhanced production in the winter period; in contrast elements of natural origin show summer concentration levels higher than the winter ones, possibly as a consequence of an increased re-suspension of soil matter in summer. The values in the two tables are quite similar for some elements whereas difference distributions between the two time periods are evident. This last issue influences the element behavior in atmosphere.

As a very simple approach for understanding the element behavior, they were grouped into three categories according to the value of the ratio (R) of summer to winter average levels (Table 6). The first group includes elements whose R is greater than 2; the second group elements whose R is less than 2 but greater than 0.5; the last group includes elements whose R is less than 0.5. Looking at the table, it can be noted a ratio difference between the elements determined in outskirts and in downtown: only 6 elements, such as Al, Br, K, Na, Ti, U, show $R > 2$ in samples collected in downtown whereas 20 elements, i.e. Al, Ba, Ca, Ce, Cs, Eu, Fe, Hf, K, La, Mg, Na, Rb, Sc, Ta, Tb, Th, Ti, U, Zr, show ratios > 2 in samples collected in outskirts.

Elements of natural origin are only found in the first group, while elements of both natural and anthropogenic origin are present in the second group. The third group includes only pollutant elements (Cd, Cr, Mo, Ni, Pb, V, Zn).

An interesting consideration, coming from the data analysis, are obviously strictly related to the general meteorological characteristics of Italy and therefore contain some peculiarity. If they are compared to similar results obtained in other countries under different meteorological conditions, it can be seen that they agree fairly well for the pollutant elements, whereas for most elements of natural origin there are sensible differences that may be related to the geomorphological and meteorological characteristics. In fact, the higher values

Table 3 Correlation coefficients for the trend of concentrations of analyzed elements present in PM_{2.5}

As	Au	Ba	Br	Ce	Co	Cr	Cs	Fe	Hf	Hg	La	Mo	Rb	Sb	Sc	Se	Sm	Th	W	Zn	
	0.25	-0.80	0.85	0.37	0.85	0.65	0.50	0.80	0.51	-0.35	0.70	0.50	0.39	0.83	0.40	0.80	0.55	0.20	-0.19	0.37	As
		-0.97	0.58	0.51	0.35	0.06	0.54	0.33	0.73	0.35	0.01	0.10	0.77	0.48	0.01	0.56	0.08	-0.32	-0.19	0.16	Au
			-0.97	-1.00	-0.98	-0.95	-0.89	-0.65	-1.00	-0.43	-0.81	-0.91	-1.00	-0.99	0.62	-0.94	-0.98	-0.98	-0.11	-0.65	Ba
				0.61	0.87	0.54	0.57	0.83	0.86	-0.22	0.57	0.23	0.68	0.95	0.35	0.89	0.50	0.16	-0.17	0.58	Br
					0.51	0.39	0.29	0.46	0.59	0.15	0.41	0.15	0.77	0.64	0.05	0.66	0.10	-0.32	-0.14	0.50	Ce
						0.69	0.53	0.81	0.63	-0.22	0.72	0.24	0.78	0.90	0.35	0.86	0.53	0.25	0.10	0.60	Co
							0.09	0.67	-0.13	-0.34	0.66	0.43	0.45	0.61	0.29	0.71	0.45	0.04	-0.24	0.44	Cr
								0.44	0.83	0.15	0.47	0.19	0.72	0.60	0.13	0.43	0.19	0.18	-0.20	0.05	Cs
									0.39	-0.38	0.73	0.14	0.57	0.85	0.65	0.79	0.72	0.27	-0.29	0.50	Fe
										0.56	0.13	0.48	-0.05	0.82	0.26	0.73	0.46	0.26	0.01	0.55	Hf
											-0.34	-0.16	0.22	-0.17	-0.41	-0.21	-0.40	-0.37	-0.03	-0.04	Hg
												0.30	0.49	0.64	0.42	0.60	0.55	0.41	-0.15	0.26	La
													0.36	0.27	-0.07	0.40	0.12	-0.31	-0.21	0.03	Mo
														0.79	-0.34	0.76	-0.29	-0.57	-0.02	0.37	Rb
															0.41	0.87	0.53	0.15	-0.13	0.68	Sb
																0.24	0.89	0.47	-0.11	0.44	Sc
																	0.44	-0.06	-0.28	0.53	Se
																		0.39	-0.09	0.50	Sm
																			0.24	0.17	Th
																				0.07	W
																					Zn

Table 4 Seasonal element concentrations (average, min and max levels expressed as ng m⁻³; a: pg m⁻³) in PM₁₀ determined by INAA and IPAA in atmospheric particulate sampled in outskirts of Rome

Element	Summer			Winter		
	Average	Min	Max	Average	Min	Max
Ag	0.42	0.2	0.9	0.31	0.1	0.5
Al	3800	200	14000	1800	400	8500
As	2	0.5	9	3	0.2	15
Ba	4	2	8	13	3	57
Br	14	3	51	5	0.4	18
Ca	2400	300	5800	1800	300	2500
Cd	0.9	0.2	4	0.9	0.2	5
Ce	1	0.09	3	0.7	0.1	4
Cl	1300	300	9300	1400	200	5400
Co	0.4	0.05	2.1	0.3	0.04	2.4
Cr	6	1	12	9	1	21
Cs	0.2	0.01	1.9	0.3	0.01	1.6
Eu ^a	0.15	0.02	0.22	0.11	0.02	0.24
Fe	1850	600	4800	1500	350	4100
Hf ^a	91	21	320	75	11	250
Hg	0.64	0.1	1.6	0.45	0.1	1.5
I	6	2	12	4	3	9
K	930	100	3400	490	50	2800
La ^a	12	1	38	7	0.9	25
Mg	1490	100	3500	950	100	1500
Mn	52	12	98	37	10	85
Mo	3.4	0.4	12	2.8	0.2	11
Na	2500	300	9800	1530	200	5200
Ni	5.5	1	19	4	1	15
Pb	42	12	157	54	15	250
Rb	5	1	18	6	1	24
Sb	8	0.5	17	9	0.4	31
Sc ^a	45	14	120	51	12	135
Se	0.7	0.1	1.3	0.6	0.1	1.6
Ta	0.02	0.01	0.09	0.03	0.01	1.1
Tb	0.048	0.009	0.32	0.035	0.005	0.42
Th ^a	200	22	510	180	19	460
Ti	250	50	930	124	100	630
Tl	1.2	0.3	6	1.7	0.5	9
U	0.3	0.02	0.9	0.2	0.03	0.8
V	15	1	34	21	5	49
Zn	130	25	290	180	40	640
Zr	45	5	75	39	7	61

Table 5 Seasonal element concentrations (average, min and max levels expressed as ng m⁻³) in PM₁₀ determined by INAA and IPAA in atmospheric particulate sampled in downtown Rome

Element	Summer			Winter		
	Average	Min	Max	Average	Min	Max
Ag	0.22	0.1	0.5	0.25	0.1	0.6
Al	2900	500	5300	800	200	1600
As	6	1	15	4	1	9
Ba	60	30	120	30	5	70
Br	40	20	70	70	10	140
Ca	2200	800	4200	1200	300	2000
Cd	0.4	0.3	0.9	0.75	0.3	2
Ce	5.6	0.7	10	2	0.2	5
Cl	1300	300	3900	1400	300	4700
Co	0.7	0.4	1.2	0.5	0.1	0.9
Cr	7	4	13	16	2	38
Cs	1.2	0.6	2.2	0.6	0.2	1.3
Eu	0.08	0.04	0.14	0.03	0.01	0.07
Fe	2200	1000	3600	1100	400	2700
Hf	0.36	0.15	0.62	0.13	0.01	0.30
Hg	0.14	0.05	0.36	0.13	0.03	0.20
I	6	3	8	6	3	10
K	1120	50	1900	320	50	1000
La	5	2	9	2	0.3	4
Mg	1090	50	2700	560	200	1200
Mn	40	15	60	35	10	90
Mo	0.5	0.2	0.6	2	0.3	7
Na	1500	300	7000	730	200	3500
Ni	1.7	1	5	8	3	13
Pb	120	80	200	270	80	840
Rb	17	10	30	7	2	15
Sb	2	0.7	4	2	0.7	7
Sc	0.3	0.1	0.5	0.1	0.03	0.2
Se	0.7	0.2	1.1	0.5	0.2	1.3
Ta	0.03	0.01	0.05	0.01	0.005	0.02
Tb	0.055	0.1	0.10	0.024	0.01	0.05
Th	3	1	5	0.9	0.3	2.5
Ti	540	200	1050	265	200	500
Tl	1	0.5	2	1.4	0.5	4
U	0.4	0.1	0.7	0.2	0.06	0.4
V	8	1	15	14	8	23
Zn	110	50	260	190	50	520
Zr	50	10	90	20	5	50

Table 6 Grouping of elements in PM₁₀ according to the ratio of summer to winter seasonal average

Ratio >2	Ratio ~1	Ratio <1
<u>Outskirt</u>		
Al, Ba, Ca, Ce, Cs, Eu, Fe, Hf, K, La, Mg, Na, Rb, Sc, Ta, Tb, Th, Ti, U, Zr	Ag, As, Br, Cl, Co, Hg, I, Mn, Sb, Se, Tl	Cd, Cr, Mo, Ni, Pb, V, Zn
<u>Downtown</u>		
Al, Br, K, Na, Ti, U	Ag, Ca, Cd, Ce, Co, Eu, Fe, Hf, La, Mg, Mn, Mo, Ni, Se, Tb, Th, Zr	As, Ba, Cl, Cr, Cs, I, Pb, Rb, Sb, Sc, Ta, Ti, V, Zn

found in this study for Al, Cs, Na, Rb, Th, Ti, U and rare earths are to be related to the element content of the volcanic rocks which are very widespread in Latium [30,31].

The enrichment factor (EF) application

In order to investigate a retrospective study of elements in PM₁₀ and their evolution in relationship with the natural or anthropogenic origins, Table 7 reports the levels of selected elements collected in the last 4 decades: the data obtained show a decrease ranging between 24% (Co) and 91% (La), except for Hg, Sb and Se. Basically, this may be attributed to the technological growth during the entire period and to the adoption of anti-pollution system in domestic heating and in industrial plants. Mercury decreases slowly (during the four decades the values are almost similar ranging between 0.09 and 0.13) whereas Se increases except during the period 1989–92 (probably due to the sample site choice), and Sb is 4-times higher than the seventies. For a better knowledge of this evolution and, especially, of the element origin, we have calculated the enrichment factors

Table 7 Levels (ng m⁻³) of selected elements in PM₁₀ investigated along four decades in downtown Rome

	PM ₁₀		
	1965-78	1989-92	2000-05
As	4.00		1.35
Br	70	50	22
Cd	0.751		0.520
Co	0.498	0.523	0.379
Cr	16	2.3	7.28
Hg	0.131	0.092	1.07
La	2.04	0.803	0.188
Ni	8.01	1.72	4.54
Pb	270	172	92
Sb	1.99	2.13	9.22
Se	0.533	0.091	0.692
Th	0.911	0.723	0.229
V	14	4.82	4.02
Zn	190	28	80




Figure 1 EF trend comparison of selected trace elements in PM₁₀ fraction calculated using La as normalizing element. ▲: period 1965–78; □: period 1989–92; ◆: period 2000–05.

(EFs) with respect to the element abundance in the upper continental crust. Elements with EF values much higher than 1 can be considered of non-crustal origin and may be attributed to long-transport phenomena from other natural and/or anthropogenic sources. The EFs have been calculated in according to the equation reported in refs. [7,32] and La as normalizing element [21]. Figure 1 shows the EF trend for selected elements in PM₁₀ during four decades. As can be noted, all the elements may be attributed to long-range transport phenomena from other natural and/or anthropogenic sources: this behavior is common to all the period studied even if a very light decreasing trend can be evidenced from 1970 to 2005.

In Figure 1, showing the EFs of some elements, three groups can be identified: La and Th ranging between 1 and 5; Co, As, V, Cr and Ni between 1 and 100; Hg, Zn, Se, Pb, Sb, Cd and Br ranging between 200 and 2500.

Basically, some particular considerations can be extrapolated: the high EF values found for Br (and Pb as well) by both the elaborations could be attributed to the use of leaded gasoline (cars with leaded gasoline are still present at the end of nineties'); the sources of As, Pb, Sb and Zn would be looked for among the various anthropogenic activities in the Rome area and particularly Sb and Zn could be of traffic origin being essential components of anti-friction alloys and car tires.

Finally, a same approach has been performed to elements investigated in the PM_{2.5} fraction, even if no historical data




Figure 2 EFs of selected elements in PM_{2.5} fraction calculated using La as normalizing element.

Table 8 Nuclear data and LOD of elements by INAA and IPAA (m: minutes; h: hours; d: days; y: years)

Element	Product nuclide	Half life		γ-Ray used (keV)	LOD (g)
<i>INAA</i>					
Ag	^{110m} Ag	253.0	d	657.8	1×10 ⁻¹⁰
Al	²⁸ Al	2.3	m	1778.9	3×10 ⁻⁸
As	⁷⁶ As	26.3	h	559.2	1×10 ⁻¹²
Au	¹⁹⁸ Au	2.70	d	411.8	6×10 ⁻¹⁴
Ba	¹³¹ Ba	11.5	d	496.3	2×10 ⁻¹⁰
Br	⁸² Br	1.47	d	776.5	2×10 ⁻¹²
Ca	⁴⁹ Ca	8.8	m	3083.0	2×10 ⁻⁷
Cd	^{115m} In	53.0	h	336.6	3×10 ⁻¹¹
Ce	¹⁴¹ Ce	32.38	d	145.4	8×10 ⁻¹⁰
Cl	³⁸ Cl	37.3	.	1642.0	2×10 ⁻⁹
Co	⁶⁰ Co	5.272	y	1332.5	6×10 ⁻¹¹
Cr	⁵¹ Cr	27.7	d	320.0	1×10 ⁻¹⁰
Cs	¹³⁴ Cs	2.062	y	795.7	1×10 ⁻¹¹
Eu	¹⁵² Eu	12.7	y	1408.0	2×10 ⁻¹²
Fe	⁵⁹ Fe	45.1	d	1099.2	2×10 ⁻⁸
Hf	¹⁸¹ Hf	42.4	d	482.2	2×10 ⁻¹¹
Hg	²⁰³ Hg	46.9	d	279.0	4×10 ⁻¹¹
I	¹²⁸ I	25.0	m	442.7	8×10 ⁻¹¹
K	⁴² K	12.52	h	1524.7	2×10 ⁻¹⁰
La	¹⁴⁰ La	40.27	h	1596.2	1×10 ⁻¹²
Mg	²⁷ Mg	9.5	m	1014.1	2×10 ⁻⁸
Mn	⁵⁶ Mn	2.58	h	1810.7	8×10 ⁻¹³
Mo	⁹⁹ Mo	2.75	d	141.0	3×10 ⁻¹¹
Na	²⁴ Na	15.0	h	1368.4	4×10 ⁻¹²
Nd	¹⁴⁷ Nd	11.1	d	531.0	1×10 ⁻¹¹
Ni	⁵⁸ Co	70.78	d	810.7	6×10 ⁻⁹
Rb	⁸⁶ Rb	18.66	d	1076.7	8×10 ⁻¹⁰
Sb	¹²² Sb	2.70	d	564.0	1×10 ⁻¹²
Sc	⁴⁶ Sc	83.85	d	889.2	2×10 ⁻¹²
Se	⁷⁵ Se	120.4	d	264.6	1×10 ⁻¹⁰
Sm	¹⁵³ Sm	1.948	d	103.1	6×10 ⁻¹³
Ta	¹⁸² Ta	115	d	1221.6	1×10 ⁻¹¹
Tb	¹⁶⁰ Tb	72.1	d	879.4	2×10 ⁻¹²
Th	²³³ Pa	27.4	d	311.8	1×10 ⁻¹¹
Ti	⁵¹ Ti	5.8	m	320.0	1×10 ⁻⁸
U	²³⁹ Np	2.35	d	228.2	5×10 ⁻¹²
V	⁵² V	3.76	m	1434.4	9×10 ⁻¹⁰
W	¹⁸⁷ W	24.0	h	685.7	1×10 ⁻¹²
Yb	¹⁷⁵ Yb	4.21	d	396.1	1×10 ⁻¹²
Zn	⁶⁵ Zn	243.8	d	1115.5	1×10 ⁻¹⁰
<i>IPAA</i>					
As	⁷⁶ As	1.097	d	559.1	10 ⁻⁷

Table 8 Nuclear data and LOD of elements by INAA and IPAA (m: minutes; h: hours; d: days; y: years) (Continued)

Ca	⁴⁷ Ca	4.536	d	1297.1	10 ⁻⁷
Ce	¹⁴¹ Ce	32.5	d	145.4	10 ⁻⁷
Cr	⁵¹ Cr	27.7	d	320.0	10 ⁻⁸
Cs	¹³⁴ Cs	2.062	y	795.9	10 ⁻⁸
Mn	⁵⁴ Mn	312.2	d	834.8	10 ⁻⁶
Nb	^{92m} Nb	10.15	d	934.5	10 ⁻⁸
Pb	²⁰⁴ Pb	52.1	h	279.0	10 ⁻⁷
Sr	⁸⁵ Sr	64.84	d	514.0	10 ⁻⁶
Ti	⁴⁷ Sc	3.341	d	159.4	10 ⁻⁸
Tl	²⁰³ Tl	12.0	d	440.0	10 ⁻⁷
Y	⁸⁸ Y	106.61	d	1836.1	10 ⁻⁷
Zr	⁹⁰ Zr	79.4	h	909.0	10 ⁻⁸

LODs calculated according to ref. [36] for matrix free elements, i.e. without disturbing activities from a sample matrix.

series are available. Figure 2 reports the results obtained on the PM_{2.5} fraction: Co, Cr, As, Zn, Hg, Sb, Se and Br show EF values ranging between 5 and 8500, respectively.

It should be noted that the EFs in PM_{2.5} fraction is more elevated that in PM₁₀ fraction, especially for Br, Se, Sb and Hg: this could be due to the different granulometric size and the different ability to penetrate of such elements. This occurrence can be an index of the different bioavailability of an element series present in PM_{2.5} fraction compared to PM₁₀ fraction. As reported above, the higher EF value found for Br and attributed to the use of leaded, is more evident in this fraction.

Conclusions

The experimentation has been addressed for getting the maximum analytical informative ability from the single sample determinations. The INAA and IPAA techniques allow to reach such elevated sensibility/accuracy levels to furnish discreet values for elements present at very low concentrations (trace and ultra-trace levels). In particular, IPAA is a unusual technique but its coupling with INAA, another nuclear methodology, can be useful for solving a lot of analytical problems.

The element concentrations determined in this study do not seem to show significant level of attention from a toxicological point of view, stressing preliminary results obtained in previous studies [7,9]. On the other hand, the element behavior trend during these last 40 years in downtown Rome is lightly decreasing although their main source is the anthropogenic activities.

Experimental Sampling

The sample collection has been performed in a large period of time, almost 40 years: this means that the operation sampling has been carried out by means of different

samplers. During the seventies the sampling occurred with no cut-off whereas since nineties the sampling heads have been introduced allowing to separate different granulometric fraction (firstly PM₁₀ and after PM_{2.5}). In particular, the latter campaigns were performed by a dichotomous sampler (mod. SA 241, Graseby,-Andersen) operating at 16.7 L min⁻¹. This sampler has a PM10 size selective inlet and separates the aerosol into fine (aerodynamic diameter, D_a, <2.5 μm) and coarse (2.5 μm < D_a < 10 μm) fractions. Particulate matter was collected on polymethylpentane ringed, 2.0 μm pore size, 37 μm, Teflon membranes (Gelman, type R2PJ). This sampler has been designed as reference by USA-EPA. The PM_{2.5} samples were stored in box at controlled conditions (atmosphere and temperature).

About 300 air samples have been collected in downtown Rome (via Urbana, an area characterized by high presence of anthropogenic activities) and Roman outskirts (Anguillara area, a rural site); all the sampling were performed at ground level. The sampling was 24-hr long for each filter. All the storage and handling sample treatment were carried out at the ENEA and ISPESL laboratories.

INAA and IPAA analyses

Among the different analytical methodologies available for element determination we used nuclear approach for its important analytical properties [7,9,20,21,33]. In fact, the various analytical techniques (spectroscopy, electrochemistry, chromatography, bioanalytical methods) [34,35] do not permit to have the maximum information because of their limitations. INAA is well known as reference analytical technique because all the experimental steps are totally traceable and there is absence of physical-chemical sample manipulation reducing the positive and/or negative artifacts formation [7,9,20,21]. Furthermore, because of its high sensitivity, multi-elemental character allowing the determination of about 40 elements with a good Limit of





Figure 3 Triga nuclear reactor (1 MW) core with Cherenkov effect.



Detection (LOD) [36] (Table 8) and accuracy, the INAA has surpassed other instrumental methods for trace/ultra-trace metal and rare earths analysis: a comparative study [37-41], however, has pointed out that INAA is blank free and especially suitable for the analysis of reference materials [41]. Further, we used IPAA as a complementary technique for determining elements: in particular, Pb, Tl and Zr which are difficult to determine by INAA, are important from toxicological and environmental point of view.

Table 8 shows the nuclear data (as product nuclide, half-life and energy peak emission) and LOD (expressed as ng m⁻³) of each element investigated in this study by means of INAA and IPAA. It should be noted the very low LODs reached by nuclear techniques in relation with other analytical methodologies [20,22].

INAA - Samples, blank and standards, put in nuclear-grade polyethylene cylinders (Kartell), were irradiated at a neutron flux of 2.6×10¹² n×cm⁻²×s⁻¹ for 32.55 h in rotatory rack "Lazy Susan" of the nuclear reactor Triga Mark II of the ENEA-Casaccia Laboratories (Figure 3) [42].

For the analysis primary and secondary standards were used. Primary standards (Carlo Erba, Milano, Italy) were As, Cd, Co, Cr, Cs, Fe, Hg, La, Ni, Sb, Se, Sm and Zn whereas as secondary standards United States Geochemical Survey (U.S.G.S.) nn. 1, 4, 6 and the Coal Fly Ash (NIST) n. 1633a were used.

After irradiation, γ-ray spectrometry measurements of different duration (Figure 4) were carried out using a HPGe detector (FWHM 1.68 keV at 1332 keV) connected to a multichannel analyzer equipped with software packages for a γ-spectra analysis.

A first measurement series was performed 5/7 days after the end of irradiation with measurement times of 3000 and 9000 s for sample for determining As, Au, Br, Cd, La, Mo, Sb, Sm, W [7,13,40]. The second series was performed 20/120 days after the end of irradiation with measurement times of 24-100 h for sample for determining Ce, Co, Cr, Cs, Eu, Fe, Hf, Hg, Nd, Ni, Rb, Sb, Sc, Se, Ta, Th, Yb, Zn [7,25,40].

IPAA - Samples, blank and standards (NIST SRM 1571) were irradiated in the photon beam of the INFN Frascati National Laboratory Linear Accelerator (LINAC) at an average beam current of 40 μA, maximum electron energy of 300 MeV and a W converter of 0.3 mm thickness.

Two series of measurements were carried out: after 36/70 hours As, Ca, Pb, Ti and Zr were measured for 2 hours whereas after 20 days from irradiation Ce, Cr, Cs, Mn, Nb, Sr, Tl and Y were counted for 4 hours.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

PA, MVR and GC coordinated the study. PA and AR set up the analytical procedure using INAA. MM processed data and provided the comparison with other literature. PA and AR edited the text and prepared the final draft of the paper. GC and MVR approved the final version. All the authors have read and approved the final manuscript.

Acknowledgements

This work was supported by INAIL under grants P20L09 and P20L01.

Author details

¹DIPIA, INAIL settore Ricerca, Certificazione e Verifica, Via IV Novembre 144, Rome 00187, Italy. ²UTFISS CATNUC, ENEA, Via Anguillarese 301, Rome 00123, Italy. ³Department of Environment, Food and Agriculture, University of Molise, Via De Sanctis, Campobasso 86100, Italy.

Received: 4 April 2013 Accepted: 24 October 2013

Published: 6 November 2013

References

1. Dockery DW, Speizer FE, Stram DO, Ware JH, Spengler JD, Ferris BG Jr: **Effects of inhalable particles on respiratory health of children.** *Am Rev Respir Dis* 1989, **139**(3):587–594.
2. Dockery DW, Pope CA: **Acute respiratory effects of particulate air pollution.** *Annu Rev Public Health* 1994, **15**:107–132.
3. Harrison RM, Yin J: **Particulate matter in the atmosphere: which particle properties are important for its effects on health?** *Sci Total Environ* 2000, **249**(1–3):85–101.
4. Cao L, Tian W, Ni B, Zhang Y, Wang P: **Preliminary study of airborne particulate matter in a Beijing sampling station by instrumental neutron activation analysis.** *Atmos Environ* 2002, **36**(12):1951–1956.
5. Avino P, Brocco D, Cecinato A, Lepore L, Balducci C: **Carbonaceous components in atmospheric aerosol: measurements, procedures and characterization.** *Ann Chim (Rome)* 2002, **92**(4):333–341.
6. Avino P, Brocco D: **Carbonaceous aerosol in the breathable particulate matter (PM10) in urban area.** *Ann Chim (Rome)* 2004, **94**(9–10):647–653.
7. Avino P, Capannesi G, Rosada A: **Characterization and distribution of mineral content in fine and coarse airborne particle fractions by neutron activation analysis.** *Toxicol Environ Chem* 2006, **88**(4):633–647.
8. Avino P, Manigrasso M: **Vertical distribution of carbonaceous material in urban atmosphere.** *Fresenius Environ Bull* 2006, **15**(8b):866–877.
9. Avino P, Capannesi G, Rosada A: **Heavy metal determination in atmospheric particulate matter by Instrumental Neutron Activation Analysis.** *Microchem J* 2008, **88**(2):97–106.
10. Manigrasso M, Febo A, Guglielmi F, Ciambottini V, Avino P: **Relevance of aerosol size spectrum analysis as support to qualitative source apportionment studies.** *Environ Pollut* 2012, **170**:43–51.
11. Manigrasso M, Avino P: **Fast evolution of urban ultrafine particles: implications for deposition doses in the human respiratory system.** *Atmos Environ* 2012, **51**:116–123.
12. Agency IAE: *Analytical Applications of Nuclear Techniques.* Wien: IAEA; 2004. ISBN 9201147031.
13. Avino P, Santoro E, Sarto F, Violante V, Rosada A: **Neutron activation analysis for investigating purity grade of copper, nickel and palladium thin films used in cold fusion experiments.** *J Radioanal Nucl Chem* 2011, **290**(2):427–436.
14. Capannesi G, Rosada A, Manigrasso M, Avino P: **Rare earth elements, thorium and uranium in ores of the North- Latium (Italy).** *J Radioanal Nucl Chem* 2012, **291**(1):163–168.
15. Segebade C, Weise HP, Lutz GJ: *Photon Activation Analysis.* Berlin: Walter de Gruyter Inc; 1988. ISBN: 0899253059.
16. Oprea C, Maslov OD, Gustova MV, Oprea IA, Mihul A, Belov AG, Szalanski PJ, Buzguta V: **Photoneutron activation analysis applied for environmental researches.** *Rom Rep Phys* 2011, **63**(2):348–356.
17. Mizera J, Randa Z: **Instrumental neutron and photon activation analyses of selected geochemical reference materials.** *J Radioanal Nucl Chem* 2010, **284**(1):157–163.
18. Campanella L, Crescentini G, Avino P, Moauro A: **Determination of macrominerals and trace elements in the alga *Spirulina platensis*.** *Analisis* 1998, **26**:210–214.
19. Avino P, Carconi PL, Lepore L, Moauro A: **Nutritional and environmental properties of algal products used in healthy diet by INAA and ICP-AES.** *J Radioanal Nucl Chem* 2000, **244**(1):247–252.
20. Capannesi G, Diaco L, Rosada A, Avino P: **Investigation of trace and ultra-trace elements of nutritional and toxicological significance in Italian potable waters by INAA.** *J Radioanal Nucl Chem* 2008, **278**(2):353–357.
21. Capannesi G, Rosada A, Avino P: **Elemental characterization of impurities at trace and ultra-trace levels in metallurgical lead samples by INAA.** *Microchem J* 2009, **93**(2):188–194.
22. Avino P, Capannesi G, Diaco L, Rosada A: **Multivariate analysis applied to trace and ultra-trace elements in Italian potable waters determined by INAA.** *Curr Anal Chem* 2010, **6**(1):26–36.
23. Capannesi G, Rosada A, Avino P: **Radiochemical separation and anti-Compton analysis of Ni, Sn, Te and Zn in lead standard reference materials at ultra-trace levels.** *Curr Anal Chem* 2010, **6**(3):217–222.
24. Buonanno G, Stabile L, Avino P, Vanoli R: **Dimensional and chemical characterization of particles at a downwind receptor site of a waste-to-energy plant.** *Waste Manag* 2010, **30**(7):1325–1333.
25. Avino P, Capannesi G, Rosada A: **Ultra-trace nutritional and toxicological elements in Rome and Florence drinking waters determined by Instrumental Neutron Activation Analysis.** *Microchem J* 2011, **97**(2):144–153.
26. Avino P, Capannesi G, Manigrasso M, Sabbioni E, Rosada A: **Element assessment in whole blood, serum and urine of three Italian healthy sub-populations by INAA.** *Microchem J* 2011, **99**(2):548–555.
27. Avino P, Capannesi G, Renzi L, Rosada A: **Instrumental neutron activation analysis and statistical approach for determining baseline values of essential and toxic elements in hairs of high school students.** *Ecotoxicol Environ Saf* 2013, **92**:206–214.
28. Randa Z, Frána J, Mizera J, Kucera J, Novák JK, Ulrych J, Belov AG, Maslov OD: **Instrumental neutron and photon activation analysis in the geochemical study of phonolitic and trachytic rocks.** *Geostand Geoanal Res* 2007, **31**(3):275–283.
29. Avino P, Capannesi G, Lopez F, Rosada A: **Determination of interesting toxicological elements in PM_{2.5} by neutron and photon activation analysis.** *Sci World J* 2013, article ID 458793. doi: 10.1155/2013/458793.
30. Locardi E, Sircana S: **Distribuzione dell'uranio e del torio nelle vulcaniti quaternarie alcaline del Lazio settentrionale.** *Rend Soc Mineralog Ital* 1967, **23**:163–224.
31. Locardi E, Mitterpergher M: **Relationship between some trace elements and magmatic processes.** *Inter J Earth Sci (Geologische Rundschau)* 1967, **57**(1):313–334.
32. Bergamaschi L, Rizzio E, Valcuvia MG, Verza G, Profumo A, Gallorini M: **Determination of trace elements and evaluation of their enrichment factors in Himalayan lichens.** *Environ Pollut* 2002, **120**(1):137–144.
33. Campanella L, Crescentini G, Avino P: **Chemical composition and nutritional evaluation of some natural and commercial food products based on *Spirulina*.** *Analisis* 1999, **27**(6):533–540.
34. Bruni F, Manzari C, Filice M, Polosa PL, Colella M, Carmone C, Hambardjjeva E, Garcia-Diaz M, Cantatore P, Roberti M: **D-MTERF5 is a novel factor modulating transcription in *Drosophila* mitochondria.** *Mitochondrion* 2012, **12**(5):492–499.
35. Cardone A, Lopez F, Affortunato F, Busco G, Hofer AM, Mallamaci R, Martinelli C, Colella M, Farinola GM: **An aryleneethynylene fluorophore for cell membrane staining.** *BBA-Biomembranes* 2012, **1818**(11):2808–2817.
36. Currie LA: **Limits for qualitative detection and quantitative determination. Application to radiochemistry.** *Anal Chem* 1968, **40**(3):586–593.
37. Farinha MM, Freitas MC, Almeida SM, Reis MA: **Some improvements in air particulate matter analysis by INAA.** *Radiat Phys Chem* 2001, **61**(3):659–661.
38. Carmo Freitas M, Almeida SM, Reis MA, Oliveira OR: **Monitoring trace elements by nuclear techniques in PM10 and PM2.5.** *Nucl Instrum Methods Phys Res A* 2003, **505**(1–2):430–434.

39. Almeida SM, Reis MA, Freitas MC, Pio CA: **Quality assurance in elemental analysis of airborne particles.** *Nucl Instrum Methods Phys Res, Sect B* 2003, **207**(4):434–446.
40. Djingova R, Arpadjan S, Kuleff I: **INAA and flame AAS of various vegetable reference materials.** *Fresenius J Anal Chem* 1991, **339**(3):181–186.
41. Djingova R, Kuleff I: **Instrumental techniques for trace analysis.** In *Trace Elements - Their Distribution and Effects in the Environment*. Edited by Markert B, Friese K. Amsterdam, The Netherlands: Elsevier; 2000:137–185. ISBN ISBN-13: 9780444505323.
42. Di Palo L, Focaccia G, Lo Prato E, Marsili F, Paletti A, Perni M, Prospero D, Sciuti S, Verri A: **Il reattore RC-1 ad 1 MW del Centro Studi Nucleari della Casaccia. Caratteristiche generali e programmi di ricerca.** *Ener Nuc* 1967, **14**:659–664.

doi:10.1186/1752-153X-7-173

Cite this article as: Avino et al.: Deep investigation on inorganic fraction of atmospheric PM in Mediterranean area by neutron and photon activation analysis. *Chemistry Central Journal* 2013 **7**:173.

Publish with **ChemistryCentral** and every scientist can read your work free of charge

“Open access provides opportunities to our colleagues in other parts of the globe, by allowing anyone to view the content free of charge.”

W. Jeffery Hurst, The Hershey Company.

- available free of charge to the entire scientific community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
<http://www.chemistrycentral.com/manuscript/>



ChemistryCentral