

RESEARCH ARTICLE

Open Access



# Tyrosinase inhibitory activity of flavonoids from *Artocarpus heterophyllous*

Hai Xuan Nguyen<sup>1</sup>, Nhan Trung Nguyen<sup>1,2</sup>, Mai Ha Khoa Nguyen<sup>1</sup>, Tho Huu Le<sup>1</sup>, Truong Nhat Van Do<sup>1</sup>, Tran Manh Hung<sup>3\*</sup> and Mai Thanh Thi Nguyen<sup>1,2\*</sup>

## Abstract

**Background:** Tyrosinase is an oxidoreductase that is very important in medicine and cosmetics because the excessive production of melanin causes hyperpigmentation. The development of novel, effective tyrosinase inhibitors has long been pursued. In preliminary tests, we found that an extract of the wood of *Artocarpus heterophyllous* (AH) potentially inhibited tyrosinase activity.

**Results:** Two new flavonoids, artocaepin E (**1**) and artocaepin F (**2**), were isolated from the wood of AH, together with norartocarpetin (**3**), artocarpanone (**4**), liquiritigenin (**5**), steppogenin (**6**), and dihydromorin (**7**). Their structures were elucidated using one-dimensional (1D) and two-dimensional (2D) nuclear magnetic resonance (NMR) and mass spectrometry. The absolute configuration of **2** was determined from the circular dichroism (CD) spectrum. Artocarpanone (**4**) had the most potent tyrosinase inhibitory effect, with an  $IC_{50}$  of  $2.0 \pm 0.1 \mu\text{M}$ , followed by artocaepin E (**1**) and steppogenin (**6**), with  $IC_{50}$  values of  $6.7 \pm 0.8$  and  $7.5 \pm 0.5 \mu\text{M}$ , respectively. A kinetic investigation indicated that **1** showed competitive inhibition, with an inhibition constant ( $K_i$ ) of  $6.23 \mu\text{M}$ .

**Conclusions:** These results demonstrate that extracts of the wood of AH and its phytochemical constituents are potential sources for skin-whitening agents.

**Keywords:** *Artocarpus heterophyllous*, Flavonoids, Chalcones, Tyrosinase inhibitors

## Background

Tyrosinase is a key enzyme in mammalian melanin synthesis. It catalyzes the first step in two reactions of melanin synthesis: the hydroxylation of L-tyrosine to 3,4-dihydroxyphenylalanine (L-DOPA) and the oxidation of L-DOPA to dopaquinone. This *o*-quinone is a highly reactive compound that can polymerize spontaneously to form melanin [1]. In mammals, melanin protects the skin from ultraviolet (UV) damage by absorbing UV sunlight and removing reactive oxygen species. However, the production of abnormal melanin pigmentation is an esthetic problem in humans [2]. Therefore, the development of high-performance tyrosinase inhibitors is much needed.

*Artocarpus heterophyllous* Lam belongs to the family Moraceae and is popularly known as jackfruit; it is distributed widely in tropical and subtropical regions of Asia. In Vietnam, this plant is known as “Mit,” and is cultivated for its edible fruits, while the wood has been used for its anti-inflammatory, antioxidant, and antiaging effects [3]. In preliminary tests, we found that an extract of the wood of AH strongly inhibited tyrosinase activity. This plant is a rich source of prenylated flavonoids and their derivatives. Numerous compounds have been isolated from AH, including prenylated flavonoids, chalcones, and simple polyphenols. These possess various biological activities, including cytotoxic, tyrosinase inhibitory, anti-inflammatory, and antioxidant effects [4–6]. The present phytochemical investigation of the wood of AH led to the isolation of two new flavonoids (**1**, **2**) and five known compounds (**3**–**7**). In this paper, we report the isolation and structure elucidation of these

\*Correspondence: hung.tran@vnu.edu.vn; nttmai@hcmus.edu.vn

<sup>2</sup> Cancer Research Laboratory, Vietnam National University, 227 Nguyen Van Cu, District 5, Ho Chi Minh, Vietnam

<sup>3</sup> Department of Biomedical Sciences, Institute for Research and Executive Education (VNUK), The University of Danang, 41 Le Duan, Haichau District, Danang, Vietnam

Full list of author information is available at the end of the article

isolated compounds, as well as their tyrosinase inhibitory activities.

## Results and discussion


### Chemistry

Dried *A. heterophyllum* wood was extracted in methanol, and the obtained extract was successively partitioned into *n*-hexane, chloroform (CHCl<sub>3</sub>), ethyl acetate (EtOAc), butanol (*n*-BuOH), and water. Repeated silica gel and reverse-phase column chromatography of the CHCl<sub>3</sub> fraction afforded two new flavonoids (1–2) and five known ones (3–7). The chemical structures of the known compounds were determined based on <sup>1</sup>H- and <sup>13</sup>C-NMR analyses and confirmed by comparison with reported spectra as norartocarpetin (3) [7], artocarpone (4) [8], liquiritigenin (5) [9], steppogenin (6) [7], and dihydromorin (7) [10] (Fig. 1).

Compound 1 was obtained as a yellow amorphous solid. High-resolution electrospray ionization mass spectrometry (HR-ESI-MS) gave a pseudo-molecular peak at *m/z* 517.1487 [M + H]<sup>+</sup> (calcd. for 517.1499), consistent with the molecular formula C<sub>29</sub>H<sub>24</sub>O<sub>9</sub>. The infrared (IR) spectrum suggested the presence of hydroxy (3410 cm<sup>-1</sup>), phenyl (1600, 1455 cm<sup>-1</sup>), and carbonyl (1710, 1700 cm<sup>-1</sup>) groups. The <sup>1</sup>H NMR spectrum of 1 included signals due to two sets of *ortho*-coupled aromatic protons at δ<sub>H</sub> 7.53 (2H, d, *J* = 8.7 Hz, H-2''' and 6''') and 6.78 (2H, d, *J* = 8.7 Hz, H-3''' and 5'''), three aromatic protons of the typical *ABX* system at δ<sub>H</sub> 6.49 (1H, d, *J* = 2.4 Hz, H-3'), 7.73 (1H, d, *J* = 8.8 Hz, H-6'), and 6.44 (1H, dd,

*J* = 8.8, 2.4 Hz, H-5'), an isolated aromatic proton at δ<sub>H</sub> 6.51 (1H, s, H-8), two *trans* olefinic protons at δ<sub>H</sub> 7.54 (1H, d, *J* = 16.0 Hz, H-7''') and 6.39 (1H, d, *J* = 16.0 Hz, H-8'''), and two isolated olefinic protons at δ<sub>H</sub> 6.98 (1H, s, H-3) and 5.55 (1H, t, *J* = 7.2 Hz, H-2''), together with one methyl, one methylene, one oxymethylene, and a characteristic signal of a hydrogen-bonded hydroxyl proton at δ<sub>H</sub> 13.34 (1H, s, 5-OH) (Table 1). In comparison, the [13] C NMR and distortionless enhancement by polarization transfer (DEPT) spectrum of 1 contained 29 carbon signals, including a ketone carbonyl, an ester carbonyl, six olefinic, a methyl, a methylene, an oxymethylene, and 18 aromatic carbons (Additional file 1). The data were similar to that of artocarpin D, which was isolated from the same plant [11], except for the presence of signals of a set of resonances due to a *trans-p*-coumaroyl unit, which was confirmed by the <sup>1</sup>H–<sup>1</sup>H correlation spectroscopy (COSY) and heteronuclear multiple bond correlation (HMBC) spectra (Fig. 2). This moiety was located at C-4'' by the HMBC correlation of H-4'' (δ<sub>H</sub> 4.51, 2H, s) with the C-9''' (δ<sub>C</sub> 166.5) ester carbonyl carbon of the *trans-p*-coumaroyl group. The nuclear Overhauser effect spectroscopy (NOESY) correlations of H-1'' with H-5'' and of H-2'' with H-4'' indicated an *E*-configuration for the C-2''/C-3'' double bond of 1. Therefore, the structure of artocarpin E was concluded to be 1.

Artocarpin F (2) was isolated as a yellow amorphous solid. The molecular formula of 2 was determined to be C<sub>20</sub>H<sub>20</sub>O<sub>5</sub> by HR-ESI-MS. The IR spectrum showed absorption bands of hydroxy (3400 cm<sup>-1</sup>), phenyl (1600,



**Fig. 1** Isolated compounds from *A. heterophyllum* (1–7)

**Table 1**  $^1\text{H}$  (500 MHz,  $J$  in Hz) and  $^{13}\text{C}$  (125 MHz) NMR data for **1** and **2**

Position	<b>1</b> <sup>a</sup>		<b>2</b> <sup>b</sup>	
	$\delta_{\text{H}}$	$\delta_{\text{C}}$	$\delta_{\text{H}}$	$\delta_{\text{C}}$
2		161.7	5.44 dd (12.8; 3.0)	80.4
3	6.98 s	106.8	2.70 dd (16.7; 3.0); 3.00 dd (16.7; 12.8)	44.6
4		182.0		191.0
5		158.4	7.59 d (8.6)	126.4
6		109.7	6.63 d (8.6)	110.5
7		161.7		162.4 <sup>c</sup>
8	6.51 s	93.2		115.5
9		155.3		162.1 <sup>c</sup>
10		103.3		130.9
1'		108.7		131.6
2'		158.8	7.42 d (8.6)	128.8
3'	6.49 d (2.4)	103.3	6.90 d (8.6)	116.2
4'		161.7		158.6
5'	6.44 dd (8.8; 2.4)	108.1	6.90 d (8.6)	116.2
6'	7.73 d (8.8)	129.8	7.42 d (8.6)	128.8
1''	3.30 d (7.2)	20.8	3.38 d (7.3)	22.4
2''	5.55 t (7.2)	126.4	5.49 t (7.3)	122.8
3''		130.2		136.3
4''	4.51 s	69.0	3.87 s	68.5
5''	1.79 s	13.9	1.65 s	13.9
1'''		125.1		
2''', 6'''	7.53 d (8.7)	130.4		
3''', 5'''	6.78 d (8.7)	115.8		
4'''		159.9		
7'''	7.54 d (16.0)	144.9		
8'''	6.39 d (16.0)	114.2		
9'''		166.5		
5-OH	13.34			

<sup>a</sup> In DMSO- $d_6$

<sup>b</sup> In acetone- $d_6$


<sup>c</sup> These signals may interchange

1450  $\text{cm}^{-1}$ ), and ketone (1705  $\text{cm}^{-1}$ ) groups. The  $^1\text{H}$  NMR spectra of **2** (Table 1) showed signals for two sets of *ortho*-coupled aromatic protons at  $\delta_{\text{H}}$  7.42 (2H, d,  $J = 8.6$  Hz,

H-2', and H-6') and 6.90 (2H, d,  $J = 8.6$  Hz, H-3' and H-5'), and others at  $\delta$  7.59 (1H, d,  $J = 8.6$  Hz, H-5) and 6.63 (1H, d,  $J = 8.6$  Hz, H-6), together with two methylene signals at  $\delta_{\text{H}}$  2.70 (1H, dd,  $J = 16.7, 3.0$  Hz, H-3a) and 3.00 (1H, dd,  $J = 16.7, 12.8$  Hz, H-3b), and one oxymethine at  $\delta_{\text{H}}$  5.44 (1H, dd,  $J = 12.8, 3.0$  Hz, H-2), which are typical of the flavanone skeleton [9]. The  $^{13}\text{C}$  NMR and DEPT spectrum of **2** displayed 20 carbon signals, including 15 carbon signals due to the flavanone skeleton and five belonging to a 4-hydroxyphenyl moiety (Fig. 1). The 4-hydroxyphenyl group was located at C-8 based on the HMBC correlations between H-1'' ( $\delta_{\text{H}}$  3.38, 2H, d,  $J = 7.3$  Hz) and C-7, C-8, C-8a, and of H-2'' ( $\delta_{\text{H}}$  5.49, 1H, t,  $J = 7.3$  Hz) with C-8 (Fig. 2). The NOESY correlations of H-1'' with H-5'' and of H-2'' with H-4'' indicated that the configuration of the C-2''/C-3'' double bond of **2** was the same as that of **1**. Finally, the absolute configuration at C-2 was considered to be *S* according to the results of the CD spectroscopic analysis, which showed negative and positive Cotton effects at 290 and 334 nm, respectively [12]. From this spectroscopic evidence, the structure of artocae-pin F was concluded to be **2**.

#### Biological assay

The tyrosinase inhibitory activity of all isolated compounds (**1**–**7**) was tested [11]. Kojic acid, a well-known tyrosinase inhibitor currently used as a cosmetic skin-whitening agent, was used as a positive control. Of the tested compounds, artocarpanone (**4**) had the most potent inhibitory effect against tyrosinase, with an  $\text{IC}_{50}$  of  $2.0 \pm 0.1$   $\mu\text{M}$ , followed by artocae-pin E (**1**) and steppogenin (**6**), with  $\text{IC}_{50}$  values of  $6.7 \pm 0.1$  and  $7.5 \pm 0.5$   $\mu\text{M}$ , respectively (Table 2). Liquiritigenin (**5**) also showed significant concentration-dependent inhibition, with an  $\text{IC}_{50}$  of  $22.0 \pm 2.5$   $\mu\text{M}$ ; this compound showed moderate inhibitory activity compared to the above compounds. However, it showed more potent inhibitory activity than kojic acid, which inhibited tyrosinase with an  $\text{IC}_{50}$  of  $44.6 \pm 0.4$   $\mu\text{M}$  (Table 2). The other compounds, artocae-pin F (**2**), norartocarpetin (**3**), and dihydromorin (**7**), showed very weak inhibitory activity, with  $\text{IC}_{50}$  values over 50  $\mu\text{M}$ .



**Fig. 2** Selected key HMBC and  $^1\text{H}$ - $^1\text{H}$  COSY correlations for **1** and **2**

**Table 2 Tyrosinase inhibitory activity of the isolated compounds 1–7**


Compounds	IC <sub>50</sub> (μM) <sup>a</sup>
<b>1</b>	6.7 ± 0.8
<b>2</b>	>50
<b>3</b>	>50
<b>4</b>	2.0 ± 0.1
<b>5</b>	22.0 ± 2.5
<b>6</b>	7.5 ± 0.5
<b>7</b>	>50
Kojic acid <sup>b</sup>	44.6 ± 0.4

<sup>a</sup> The assay was executed in triplicate

<sup>b</sup> Positive control used for enzymatic inhibition assay

Further study examined the inhibitory mechanism of artocaepin E (**1**), which strongly inhibited tyrosinase activity. To determine the type of enzyme inhibition and the inhibition constant for an enzyme-inhibitor complex, the mechanism was analyzed by Lineweaver–Burk plots. The results indicated that **1** displayed competitive inhibition, with an inhibition constant ( $K_i$ ) of 6.23 μM (Fig. 3).

On close inspection of the inhibitory activity exerted by these compounds, the following biological profile of the structure–activity relationship was deduced. In terms of the flavone skeleton, compounds **1** and **3** are derivative of apigenin, a common flavone in plants; however, the presence of one hydroxyl group at C-2', a *trans-p*-coumaroyl unit connected to the hydroxyphenyl through an ester linkage at C-6 of the apigenin skeleton in **1**, led to significantly stronger inhibitory activity than that of **3** (6.7 vs. >50 μM). This suggests that the absence of the side-chain at C-6 of the B-ring leads to a significant loss of activity, and the presence of a side-chain such as *trans-p*-coumaroyl connected to the hydroxyphenyl may positively influence the tyrosinase inhibitory activity. Regarding the flavanone skeleton, artocarpanone (**4**), which possesses a methoxyl group at C-7 of ring A, had the strongest inhibitory activity. Steppogenin (**6**) shares the same structure as **4**, except for the hydroxyl group at C-7; however, it had 3.75-fold higher inhibitory activity than **4**. In comparison, dihydromorin (**7**), which has four hydroxyl groups at C-2', C-3, C-5, and C-7, had weak activity. These results imply that the methoxyl and hydroxyl groups in the main flavanone skeleton play an important role in tyrosinase inhibition.



**Fig. 3** Lineweaver–Burk plots for type of inhibition of mushroom tyrosinase (10 U/mL) by artocaepin E (**1**) for the catalysis of L-DOPA (0.2, 0.3, 0.4, 0.5, and 0.6 mM) at 30 °C, pH 6.8. Concentration of these compounds for curves  $I_{0.0}$ ,  $I_{5.0}$ ,  $I_{7.0}$ , and  $I_{10.0}$  were 0.0, 5.0, 7.0, and 10.0 μM, respectively. The inset represents the plot of these compounds for determining the inhibition constant ( $K_i$ ). The line is drawn using a linear least squares fit

## Methods

### General procedure

Optical rotations were recorded on a JASCO DIP-140 digital polarimeter. CD measurements were carried out on a JASCO J-805 spectropolarimeter. IR spectra were measured with a Shimadzu IR-408 spectrophotometer in  $\text{CHCl}_3$  solution. NMR spectra were taken on a Bruker Advance III 500 spectrometer (Bruker Biospin) with tetramethylsilane (TMS) as an internal standard, and chemical shifts are expressed in  $\delta$  values. HR-ESI-MS measurements were carried out on a Bruker microTOF-QII spectrometer. Column chromatography was performed with BW-820MH Si gel (Fuji Silisia, Aichi, Japan). Analytical and preparative TLC was carried out on precoated Merck Kiesegel 60F<sub>254</sub> or RP-18F<sub>254</sub> plates (0.25 or 0.5 mm thickness).

### Chemicals

Tyrosinase (EC 1.14.18.1) from mushroom (3933 U/mL) and L-dihydroxyphenylalanine (L-DOPA) were obtained from Sigma Chemical Co. (St. Louis, MO, USA). Kojic acid and DMSO were purchased from Merck (Darmstadt, Germany). Other chemicals were of the highest grade available.

### Plant material

The wood of *A. heterophyllum* was collected at the Seven-Mountain area, An Giang province, Vietnam in August 2010. The plant was identified by Ms. Hoang Viet, Faculty of Biology, University of Science, Vietnam National University-Hochiminh City. The voucher sample of the wood part (AN-2985) is preserved at Department of Analytical Chemistry, Faculty of Chemistry, University of Science, Vietnam National University-Hochiminh City.

### Extraction and isolation

The dried powder of wood of *A. heterophyllum* (5.8 kg) was extracted with MeOH (15 L, reflux, 3 h,  $\times$  3) to yield a MeOH extract. The extract was partitioned between EtOAc and water to give an EtOAc-soluble fraction (64.2 g). The EtOAc-soluble fraction was subjected to silica gel column chromatography with acetone-hexane to give six fractions fr. 1–6. Fraction 6 was chromatographed further using a MeOH- $\text{CHCl}_3$  gradient system to afford four subfractions fr. 6.1–6.4. Sub-fraction 6.2 was chromatographed further using MeOH- $\text{CHCl}_3$  gradient system, with final purification effected by preparative TLC with 2 % MeOH- $\text{CHCl}_3$ , to give 4 (6.5 mg) and 5 (20.8 mg). Subfraction 6.3 was separated by preparative TLC with 5 % MeOH- $\text{CHCl}_3$  to give 1 (5.0 mg), and 2 (5.3 mg), and 3 (8.5 mg). Subfraction 6.4 was re-chromatographed on silica gel with 7 % MeOH- $\text{CHCl}_3$ , followed by final purification using preparative TLC with 40 % acetone-hexane, to give 6 (8.0 mg), and 7 (7.5 mg).

Artocaepein E (**1**): pale yellow, amorphous solid; IR  $\nu_{\text{max}}$  ( $\text{CHCl}_3$ ) 3395, 1655, 1615, 1400  $\text{cm}^{-1}$ ;  $^1\text{H}$  and  $^{13}\text{C}$  NMR (DMSO- $d_6$  500 MHz) see Table 1; HR-ESI-MS  $m/z$  517.1487 (calcd. for  $\text{C}_{29}\text{H}_{25}\text{O}_9$   $[\text{M} + \text{H}]^+$ , 517.1499).

Artocaepein F (**2**): yellowish gum;  $[\alpha]_{\text{D}}^{25} -10.0^\circ$  (c 1.0,  $\text{C}_2\text{H}_5\text{OH}$ ); IR  $\nu_{\text{max}}$  ( $\text{CHCl}_3$ ) 3365, 1630, 1600, 1510  $\text{cm}^{-1}$ ;  $^1\text{H}$  and  $^{13}\text{C}$  NMR (acetone- $d_6$  500 MHz) see Table 1; HR-ESI-MS  $m/z$  363.1224 (calcd. for  $\text{C}_{20}\text{H}_{20}\text{O}_5\text{Na}$   $[\text{M} + \text{Na}]^+$ , 363.1208).

### Tyrosinase inhibitory assay

All the samples were first dissolved in DMSO and used for the actual experiment at concentrations of  $100^{-1}$   $\mu\text{g}/\text{mL}$  (or  $\mu\text{M}$  for pure compounds). The tyrosinase inhibitory activity assay was performed as previously described by Arung et al. [13]. The assay mixtures consisting of 1900  $\mu\text{L}$  of test solution in 0.1 M phosphate buffer pH 6.8 and 100  $\mu\text{L}$  of enzyme solution (15 U/mL in 0.1 M phosphate buffer pH 6.8) was prepared immediately before use. After preincubation at room temperature for 30 min, the reaction was initiated by the addition of 1000  $\mu\text{L}$  of substrate solution (1.5 mM L-DOPA in 0.1 M phosphate buffer pH 6.8). The assay mixture was incubated at room temperature for 7 min, and the absorbance at 475 nm was measured with a Shimadzu UV-1800 spectrophotometer. Kojic acid, a known tyrosinase inhibitor, was used as positive control. Tyrosinase inhibitory activity was expressed as the percentage inhibitory of enzyme tyrosinase in the above assay system, calculated as  $(1 - B/A) \times 100$ , where  $A$  and  $B$  are the activities of the enzyme without and with test material. IC<sub>50</sub> values were calculated from the mean values of data from four determinations.

### Inhibition mechanism

The procedure for determination of the inhibition mechanism was similar to that for determination of IC<sub>50</sub>, except that uninhibited and inhibited reactions were observed for three different concentrations of L-DOPA (0.2, 0.3, 0.4, 0.5, and 0.6 mM) at 30 °C in 0.1 M phosphate buffer pH 6.8. The dependence of absorbance (475 nm) on time was measured, and the reaction rate was calculated for all reactions (uninhibited and inhibited). Then, a Lineweaver-Burk plot was constructed, and  $K_m$  and  $V_m$  values were calculated. Each measurement was performed in duplicate.

### Conclusions

In this study, we identified two new flavonoids from the wood of AH, artocaepein E (**1**) and artocaepein F (**2**), together with five known compounds: norartocarpetin (**3**), artocarpanone (**4**), liquiritigenin (**5**), steppogenin (**6**), and dihydromorin (**7**). Regarding tyrosinase inhibition, artocarpanone (**4**) had the greatest inhibitory

effect, followed by artocaepin E (1) and steppogenin (6). Liquiritigenin (5) also showed significant concentration-dependent inhibition. Kinetic studies indicated that the new active compound artocaepin E (1) displayed competitive inhibition. These results suggest that these compounds may serve as structural templates for the design and development of novel tyrosinase inhibitors as effective anti-browning agents in cosmetics.

### Additional file

**Additional file 1.** One-dimensional (1D) and two-dimensional (2D) nuclear magnetic resonance (NMR) and mass spectrometry (MS) of compounds 1-2

### Authors' contributions

HXN and NTN isolated and elucidated the compounds, MHKN and THL carried out the bioassay, TNVD and TMH wrote the manuscript, MTTN carried out conception and design of the study, read and brought some corrections to the paper. All authors read and approved the final manuscript.

### Author details

<sup>1</sup> Faculty of Chemistry, University of Science, 227 Nguyen Van Cu, District 5, Ho Chi Minh, Vietnam. <sup>2</sup> Cancer Research Laboratory, Vietnam National University, 227 Nguyen Van Cu, District 5, Ho Chi Minh, Vietnam. <sup>3</sup> Department of Bio-medical Sciences, Institute for Research and Executive Education (VNUK), The University of Danang, 41 Le Duan, Haichau District, Danang, Vietnam.

### Acknowledgements

This research is funded by Vietnam National University Hochiminh City (VNU-HCM) under Grant number A2015-18-02.

### Competing interests

The authors declare that they have no competing interests.

Received: 30 October 2015 Accepted: 10 January 2016

Published online: 29 January 2016

### References

- Seo SY, Sharma VK, Sharma N (2003) Mushroom tyrosinase: recent prospects. *J Agric Food Chem* 51:2837–2853
- Briganti S, Camera E, Picardo M (2003) Chemical and instrumental approaches to treat hyperpigmentation. *Pigment Cell Res* 16:101–110
- Vo.VC An giang medicinal plants. Sci Technol Publisher: An Giang, Vietnam. 1991.
- Septama AW, Panichayupakaranant P (2015) Antibacterial assay-guided isolation of active compounds from *Artocarpus heterophyllous* heartwoods. *Pharm Biol* 53:1608–1613
- Ruiz-Montanez G, Burgos-Hernandez A, Calderon-Santoyo M, Lopez-Saiz CM, Velazquez-Contreras CA, Navarro-Ocana A et al (2015) Screening antimutagenic and antiproliferative properties of extracts isolated from Jackfruit pulp (*Artocarpus heterophyllous* Lam). *Food Chem* 175:409–416
- Shrikanta A, Kumar A, Govindaswamy V (2015) Resveratrol content and antioxidant properties of underutilized fruits. *J Food Sci Technol* 52:383–390
- Jeong SH, Ryu YB, Curtis-Long MJ, Ryu HW, Baek YS, Kang JE et al (2009) Tyrosinase inhibitory polyphenols from roots of *Morus lhou*. *J Agric Food Chem* 57:1195–1203
- Wei BL, Weng JR, Chiu PH, Hung CF, Wang JP, Lin CN (2005) Anti-inflammatory flavonoids from *Artocarpus heterophyllous* and *Artocarpus communis*. *J Agric Food Chem* 53:3867–3871
- Kulesh NI, Vasilevskaya NA, Veselova MV, Denisenko VA, Fedoreev SA (2008) Minor polyphenols from *Maackia amurensis* wood. *Chem Nat Compd* 44:712–714
- Zheng ZP, Cheng KW, To JT, Li H, Wang M (2008) Isolation of tyrosinase inhibitors from *Artocarpus heterophyllous* and use of its extract as anti-browning agent. *Mol Nutr Food Res* 52:1530–1538
- Nguyen NT, Nguyen MHK, Nguyen HX, Bui NKN, Nguyen MTT (2012) Tyrosinase inhibitors from the wood of *Artocarpus heterophyllous*. *J Nat Prod* 75:1951–1955
- Slade D, Ferreira D, Marais JPJ (2005) Circular dichroism, a powerful tool for the assessment of absolute configuration of flavonoids. *Phytochemistry* 66:2177–2215
- Arung ET, Kusuma IW, Iskandar YM, Yasutake S, Shimizu K, Kondo R (2005) Screening of Indonesian plants for tyrosinase inhibitory activity. *J Wood Sci* 51:520–525

Submit your manuscript to a SpringerOpen® journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)